

<p>(51) International Patent Classification ⁵ : C07H 15/12, C07K 3/00, 13/00 C07K 15/00, 17/00, A61K 35/14 A61K 35/78, 35/80, 37/04 C12Q 1/68, C12N 15/00 C12P 21/00, 21/02, 21/06 C12P 19/34</p>	<p>A1</p>	<p>(11) International Publication Number: WO 93/19079</p> <p>(43) International Publication Date: 30 September 1993 (30.09.93)</p>
<p>(21) International Application Number: PCT/US93/02428</p> <p>(22) International Filing Date: 18 March 1993 (18.03.93)</p> <p>(30) Priority data: 07/855,412 19 March 1992 (19.03.92) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 855,412 (CIP) Filed on 19 March 1992 (19.03.92)</p> <p>(71) Applicant (for all designated States except US): WASHINGTON STATE UNIVERSITY RESEARCH FOUNDATION [US/US]; NE 1615 Eastgate Boulevard, Pullman, WA 99163 (US).</p>		<p>(72) Inventors; and (75) Inventors/Applicants (for US only) : RYAN, Clarence, A. [US/US]; NE 1280 Coveway, Pullman, WA 99163 (US). PEARCE, Gregory, L. [US/US]; 520 North I Street, Palouse, WA 99161 (US). McGURL, Barry, F. [US/US]; SE 635 Taylor, Apartment 3, Pullman, WA 99163 (US).</p> <p>(74) Agents: SMITH, DeAnn, F. et al.; Harness, Dickey & Pierce, P.O. Box 828, Bloomfield Hills, MI 48303 (US).</p> <p>(81) Designated States: AU, CA, JP, RU, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: SYSTEMIN</p> <div style="text-align: center; margin-top: 100px;"> <p>1 5 10 15 18</p> <p>⁻H₃N-AVQSKPPSKRDP⁺PKMQTD-COO⁻</p> </div>		
<p>(57) Abstract</p> <p>Systemin, an 18 amino acid peptide hormone and first polypeptide hormone found in plants, induces expression of defense genes in plants wounded mechanically or by predators including herbivores, insects, bacteria and viruses. The precursor for systemin is encoded as a 200 amino acid prosystemin molecule that has the systemin peptide sequence located near the carboxyl-terminus. Both a 951 bp cDNA for prosystemin and 4526 bp genomic DNA were cloned and the organization of the gene was determined. Transgenic plants constructed with antisense prosystemin DNA fail to mount a defensive response to wounding. Transgenic plants constructed with increased copy number of prosystemin genes exhibit increased resistance to wounding. Insect larval that feed on transgenic plants constructed with increased copy number of prosystemin genes exhibit decreased growth weight compared to larval that feed on wild type plants. A tomato systemin polypeptide has an amino acid sequence NH₃-AVQSKPPSKRDP⁺PKMQTD-COO⁻.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

- 1 -

SYSTEMIN**GOVERNMENT SPONSORSHIP**

This invention was made with government support under grant number DCB9104542 awarded by the National Science Foundation. The government has
5 certain rights in the invention.

RELATED APPLICATIONS

This application is a continuation-in-part of U.S. Patent Application Serial No. 07/855,412, filed March 19, 1992, which is a continuation-in-part of U.S. Patent Application Serial No. 07/528,956, filed May 25, 1990, and a continuation-in-part of
10 US/91/03685, filed May 24, 1991.

FIELD OF THE INVENTION

This invention relates to methods and materials for inducing plant defense mechanisms. More particularly, this invention relates to methods for inducing the production of plant defense proteins, such as proteinase inhibitors, and to methods
15 of regulating resistance to predators, herbivores, insect, pathogen or virus in plants by inducing or suppressing the expression of genes encoding systemin or prosystemin.

BACKGROUND OF THE INVENTION

Damage to crops by predators (*i.e.*, insects, herbivores, and pathogens, including fungi, bacteria, and viruses), results in substantial annual losses in
20 agricultural production. Man has created and employed a wide range of chemicals in attempting to reduce damage to plant crops. Many environmental problems have been created by the widespread use of chemicals that may only provide a transient level of protection for crops. Chemicals also suffer from the disadvantage that all
25 organisms in an area may be indiscriminately treated, causing needless damage to many beneficial organisms. Many chemicals are also potentially toxic to man and animals.

Attempts to reduce crop damage have included selective breeding for resistance, but resistance traits can frequently be controlled by many genes making
30 it difficult (or impossible) to genetically select a desired attribute. Decreased crop yields are also commonly encountered in resistance strains. Accordingly, there exists a strong need for compositions and processes to improve the resistance of plants under attack by herbivores.

Plants have evolved inducible defensive mechanisms that respond to attacks
35 by predators (C.A. Ryan, 1990, *Ann. Rev. Phytopathol.* 28:425; D.J. Bowles, 1990, *Ann.*

- 2 -

- Rev. Biochem*, 59:873; M. Chessin and A.E. Zipf, 1990, *The Botanical Review* 56:193; D.L. Dreyer and B.C. Campbell, 1987, *Plant, Cell and Environ.* 10:353). One mechanism involves systemic synthesis of serine proteinase inhibitors that are accumulated at distal tissue sites in plants. The proteinases can inhibit the digestive
- 5 enzymes of insects and microorganisms (T.R. Green and C.A. Ryan, 1972, *Science* 175:776; C.A. Ryan, 1978, *TIBS* 3(7):148; V.A. Hilder, A.M.R. Gatehouse, S.E. Sheerman, R.F. Barker, D. Boulter, 1987 *Nature* 330:160; R. Johnson, J. Narvaez, G.An, C.A. Ryan, 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:9871). Proteinase inhibitors can be detrimental to the growth and development of insects from a variety of genera
- 10 including *Heliothis*, *Spodoptera*, *Diabrotica* and *Tribolium* (Ryan, *supra*; Broadway, *supra*; Rechsteiner, *supra*). Several families of polypeptides have been described that inhibit serine proteinases, including: the Kunitz family (e.g., Soybean trypsin inhibitor); the Bowman-Birk family; (e.g., Soybean proteinase inhibitor); the Potato I and Potato II families; the Barley trypsin inhibitor family; and, the Squash inhibitor family.
- 15 Wounding of plants by animals, including insects, and pathogens or mechanical damage reportedly induces transcriptional activation of proteinase inhibitor genes and protein synthesis (J.S. Graham, G. Hall, G. Pearce, C.A. Ryan, 1986, *Planta* 169:399). The latter wound-response has reportedly been described in a variety of species including; tomato (J.S. Graham, G. Pearce, J. Merryweather, K. Titani, L.
- 20 Ericsson, C.A. Ryan, 1985, *J. Biol. Chem.* 260(11):6555; J.S. Graham, G. Pearce, J. Merryweather, K. Titani, L.H. Ericsson, C.A. Ryan, 1985, *J. Biol. Chem.* 260(11):6561), potato (C.A. Ryan, 1968, *Plant Physiol.* 43:1880), alfalfa (W.E. Brown and C.A. Ryan, 1984, *Biochemistry* 23:3418; W.E. Brown, K. Takio, K. Titani, C.A. Ryan, 1985, *Biochemistry* 24:2105), cucurbits (D. Roby, A. Toppan, M.T. Esquerre-Tugaye, 1987,
- 25 *Physiol. Mol. Pl. Pathol.* 30:6453) and poplar trees (H.D. Bradshaw, J.B. Hoflick, T.J. Parsons, H.R.G. Clarke, 1989, *Plant Mol. Biol.* 14:51). Wounding reportedly results in the rapid accumulation of proteinase inhibitors not only in wounded leaves but also in distal, unwounded leaves, suggesting that a signal, or signals, released from the wound site travels throughout the plant. Transport of these signals is mediated locally
- 30 through intercellular and intracellular fluids that permeate wound or infection sites (Green, T.R. and C.A. Ryan, *Science* 175:776-777, 1972) or travel systemically through the vascular system of plants (Kuc, J. and C. Presisig, *Mycologia* 76:767-784, 1984; M. Kopp, et al., *Plant Physiol.* 90:208-216, 1990; and K.E. Hammond-Kosack, et al., *Physiol. Mol Plant Path.* 35:495-506, 1989). Proposed wound signals include: pectic
- 35 fragments derived from the plant cell wall (C.A. Ryan and E.E. Fanner, 1991, *Annu.*

- 3 -

- Rev. Plant. Physiol. Mol. Bio.* 42:651); the lipid-derived molecule, jasmonic acid (E.E. Farmer and C.A. Ryan, 1990, *Proc. Natl. Acad. Sci. U.S.A.* 87:7713); the plant hormone, abscisic acid (H. Pena-Cortes, J.J. Sanchez-Serrano, R. Mertens, L. Willmitzer, S. Prat, 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:9851); electrical potentials (E. Davies, 1987, *Plant, Cell and Environ.* 10:623; J.F. Thain, H.M. Doherty, D.J. Bowles, D.C. Wildon, 1990, *Plant, Cell and Environ.* 13:569); and, more recently, an 18-amino acid polypeptide called systemin (G. Pearce, D. Strydom, S. Johnson, C.A. Ryan, 1991, *Science* 253:895).

SUMMARY OF THE INVENTION

- Disclosed herein are a) the isolation and sequencing of Systemin, an 18-amino acid polypeptide and first polypeptide hormone found in plants; b) the isolation and sequencing of prosystemin, a precursor 200 amino acid 23kDa polypeptide; c) the molecular cloning of cDNA encoding prosystemin and genomic DNA encoding prosystemin mRNA; d) the construction of antisense vectors encoding antisense RNA inhibiting prosystemin synthesis; e) the construction of vectors containing the prosystemin or systemin sense nucleic acid, as well as, f) a method of enhancing the defense mechanism of plants. Systemin has been shown to be a powerful inducer of the synthesis of wound-inducible plant defense proteins including members of proteinase inhibitor families, *i.e.*, the Inhibitor I (8100 Da) and Inhibitor II (12,300 Da) families. Radioactively labelled systemin applied to a plant wound site is rapidly translocated to distal tissues where it induces synthesis of defense proteins. Systemin is represented only once in the precursor prosystemin molecule and is located close to the carboxy terminus of the precursor protein. Plants expressing antisense prosystemin RNA exhibit a greatly reduced synthesis of wound-induced proteinase inhibitors. Transgenic plants expressing sense prosystemin RNA exhibit an increased level of wound-induced proteinase inhibitors. *Manduca sexta* larvae exhibit lowered growth weight when feeding on transgenic sense tomato plants compared to those larvae feeding on wild type plants.

- Nucleic acid sequences of the invention are capable of encoding a systemin or prosystemin polypeptide or antisense RNA. The nucleic acids comprise a nucleotide sequence capable of hybridizing with the sense or antisense strand of the nucleotide sequence of the prosystemin cDNA or genomic DNA. The nucleic acids of the invention encode prosystemin and systemin polypeptides or are antisense sequences which interfere with the expression of systemin or prosystemin *in vivo*. Systemin related polypeptides of the present invention comprise the amino acid

- 4 -

sequence $R_1R_1QR_1R_2PPR_1R_2R_2R_1PPR_2R_1QR_1R_1$, wherein R_1 is any amino acid, R_2 is lysine or arginine (or any derivative thereof), Q is glutamine (or any derivative thereof), and P is proline (or any derivative thereof). A representative example of a systemin polypeptide of the invention is the amino acid sequence: NH_3-
5 AVQSKPPSKRDPPKMQTD-COO $^-$.

The processes of the present invention are useful for enhancing synthesis of defense proteins in a plant by introducing a prosystemin sense nucleic acid into a plant cell, or inhibiting synthesis by introducing an antisense nucleic acid. Transgenic plants containing the subject nucleic acids of the invention are also provided.

10 BRIEF DESCRIPTION OF THE DRAWINGS

The foregoing aspects and many of the attendant advantages of this invention will become more readily appreciated as the same becomes better understood by reference to the following detailed description, when taken in conjunction with the accompanying drawings, wherein:

15 FIGURE 1 shows preliminary purification of systemin from the extracts of tomato leaves by semipreparative reverse-phase HPLC as described in Example 1, below.

FIGURE 2 shows substantial purification of systemin by chromatography on an SCX-HPLC column as described in Example 1, below.

20 FIGURE 3 shows the amino acid sequence of the systemin polypeptide.

FIGURE 4 shows induction of defense protein synthesis, *i.e.*, Inhibitor I (closed circles) and Inhibitor II (open circles), in tomato plants by a synthetic systemin polypeptide, as described in Example 2, below.

FIGURE 5A shows the autoradiograph of a tomato leaf that was treated with
25 ^{14}C -labeled synthetic systemin polypeptide to demonstrate transport of systemin from wound sites into distal plant tissues.

FIGURE 5B shows the ^{14}C -labeled synthetic systemin isolated by reverse-phase HPLC from the distal plant tissues of FIGURE 5A.

FIGURE 6 shows the amino acid sequence of prosystemin.

30 FIGURE 7 shows the nucleotide sequence of cDNA encoding prosystemin with crosshatched underlining showing repeated sequence motifs and vertical bar underlining showing the location of systemin in the precursor sequence.

FIGURE 8 shows the nucleotide sequence of the prosystemin gene.

FIGURE 8A shows the nucleotide sequence of the prosystemin gene from
35 position 1 to position 2100.

- 5 -

FIGURE 8B shows the nucleotide sequence of the prosystemin gene from position 2101 to position 4200.

FIGURE 8C shows the nucleotide sequence of the prosystemin gene from position 4201 to position 4526.

5 FIGURE 9A shows the organization of the prosystemin gene. The gene consists of a 104 bp 5'-untranslated region, a 4176 bp coding region composed of 11 exons (vertical bars) interrupted by 10 introns, and a 246 bp 3'-untranslated region. The position of systemin is indicated by a horizontal bar labelled SYS.

10 FIGURE 9B shows a Southern blot analysis of the prosystemin gene. Tomato genomic DNA was isolated from leaves (as described in Example 6, below), and 5 µg was digested with EcoRI (lane 1); Bgl II (lane 2) or Sca I (lane 3), and electrophoresed on a 0.8% agarose gel that was probed with nick-translated prosystemin.

15 FIGURE 9C shows a Southern blot analysis of the species distribution of prosystemin gene homologues, as described in Example 9, below. Genomic DNA (5 µg) from tomato (lane 1), potato (lane 2), tobacco (lane 3), alfalfa (lane 4), and *Arabidopsis* (lane 5) was digested with EcoRI and electrophoresed on a 0.8% agarose gel. The gel was blotted onto nitrocellulose and probed with nick-translated prosystemin cDNA.

20 FIGURE 10 shows the organization of the prosystemin gene. Exons are represented by vertical bars and numbered 1 to 11. The five exon pairs are: 1 plus 2; 3 plus 4; 5 plus 6; 7 plus 8 and 9 plus 10.

25 FIGURE 11 shows sequence alignment of the prosystemin gene exons. The consensus sequence (con) is composed of those bases that occur at the same position in at least three of the five exon sequences.

FIGURES 11A shows the alignment of sequences of the first exons of each pair (exons 1, 3, 5, 7, and 9).

FIGURES 11B shows the alignment of the sequences of the second exons of each pair (exons 2, 4, 6, 8, and 10).

30 FIGURE 12 shows sequence alignment of three repeated polypeptide sequences within prosystemin. Three polypeptide sequences (Rep A, Rep B and Rep C; each occurring once within the amino-terminal half of prosystemin), are aligned with the homologous sequences (Rep 2A, Rep 2B and Rep 2C; each occurring once within the carboxy terminal half of prosystemin). Amino acids which differ between
35 repeats are underlined. The amino acids at the beginning and end of each repeat are

- 6 -

numbered from the amino terminus of prosystemin.

FIGURE 13A shows the positions of the duplicated polypeptide sequences within prosystemin. Prosystemin is represented by a horizontal bar with the amino acid residues numbered 1 to 200 from the amino terminus. Sequence elements Rep A, Rep B and Rep C and their repeats Rep 2A, Rep 2B and Rep 2C are indicated by hatched bars. Systemin is represented by a hatched bar labelled Sys.

FIGURE 13B shows the location of the sequences encoding the polypeptide repeats within the prosystemin gene. Exons are represented by vertical bars. The parts of the exons encoding the polypeptide repeats are shaded.

FIGURE 14 shows a sequence comparison of the intron boundary at the 3'-end of the exons 3 and 7. Exon sequence is underlined. The first four bases of the intron at the 3'-end of exon 7 have been displaced to facilitate accurate alignment of the homologous sequences occurring at the 3'-end of exon 3 and at the 5'-end of the intron between exons 7 and 8.

FIGURE 15A shows a Northern blot analysis of the time course of induction of prosystemin mRNA and Inhibitor I mRNA after wounding, as described in Example 7, below.

FIGURE 15B shows a Northern blot analysis of the distribution of prosystemin mRNA in various parts of an unwounded, fully-grown tomato plant, as described in Example 8, below. Total RNA was extracted from the following parts of an unwounded tomato plant; root (R); stem (St); petiole (Pt); leaf (Le); sepal (Se); petal (Pe); stamen (Sm) and pistil (Pi).

FIGURE 16A shows a Northern blot analysis of total RNA extracted from transgenic antisense plant 1A4. Lane 1 shows the results obtained with the sense probe and Lane 2 shows the results with the antisense probe, as described in Example 10, below.

FIGURE 16B shows a graphic depiction of the levels of Inhibitor I in wounded F1 transgenic antisense plants (unshaded bars) and non-transformed control plants (solid bars), as described in Example 10, below.

FIGURE 16C shows a graphical depiction of the levels of Inhibitor II in wounded F1 transgenic antisense plants (unshaded bars) and non-transformed control plants (solid bars), as described in Example 10, below.

FIGURE 17 shows a Northern blot analysis of total RNA extracts collected at different times from undamaged leaves of control (nontransformed) and transgenic tomato plants during the feeding experiments with *Manduca sexta* larvae, as described

- 7 -

in Example 11, below.

FIGURE 18 shows the time course of accumulation of Inhibitor I and II proteins in undamaged leaves of control and transgenic tomato plants induced by the feeding of *Manduca sexta* larvae, as described in Example 11, below.

5 FIGURE 19 shows the growth of *Manduca sexta* larvae, while feeding on leaves of control and transgenic antisense tomato plants, as described in Example 11, below.

FIGURE 20 shows the time course of larval weight gain of *Manduca sexta* larvae feeding on leaves of wild type and transgenic tomato plants, as described in Example 11, below.

10 **DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT**

As used herein the following terms are used to mean:

The term "defense proteins" is intended to include proteins that impede plant tissue attack or ingestion by predators, such as by herbivores, insects, fungi, bacteria or viruses. Defense proteins increase resistance of plants to predator attack by acting
15 directly to impede plant tissue attack or ingestion, or by acting indirectly to produce other defense compounds from precursor materials, (e.g., by acting to induce enzymes in a pathway synthesizing defense compounds; or, by inducing proteins that regulate enzymes that synthesize defense compounds). Representative examples of defense proteins include: e.g., proteinase inhibitors, thionins, chitinases and β -
20 glucanases. Representative enzymes that lead to the synthesis of defense compounds include, e.g., caseinase synthase. Representative enzymes that are part of a biosynthetic pathway leading to synthesis of defense compounds includes, e.g. enzymes in the phenylpropanoid and terpenoid pathways for synthesis of phytoalexin antibiotics, alkaloids and other toxic chemicals. Other predator defense proteins
25 useful in connection with the invention disclosed herein will, of course, be apparent to those skilled in the art. Particularly suitable predator defense proteins include inhibitors of digestive proteolytic enzymes of the attacking herbivore, such as proteinase inhibitors, and antibacterial, antimycotic, and antiviral compounds and the like. Representative proteinase inhibitor defense proteins include, e.g., the Kunitz
30 family of trypsin inhibitors, the Bowman-Birk family of proteinase inhibitors, the Inhibitor I family of proteinase inhibitors, the Inhibitor II family of proteinase inhibitors, the barley family of trypsin inhibitors, and the squash family of proteinase inhibitors. Representative examples of plant proteinase inhibitors are disclosed in PCT/US/91/03685, a continuation-in-part application of U.S. Patent Application Serial
35 No. 07/528,956, the disclosures of both applications are incorporated herein by

- 8 -

reference.

The term "nucleic acid" is intended to mean natural and synthetic linear and sequential arrays of nucleotides and nucleosides, e.g., in cDNA, genomic DNA (gDNA), mRNA, and RNA, oligonucleotides, oligonucleosides, and derivatives thereof.

- 5 For ease of discussion, such nucleic acids are at times collectively referred to herein as "constructs," "plasmids," or "vectors." Representative examples of the nucleic acids of the invention include bacterial plasmid vectors such as expression, cosmid, and cloning and transformation vectors (e.g., pBR322, λ , Ti, and the like), plant viral vectors (e.g., modified CaMV and the like), and synthetic oligonucleotide molecules
10 such as chemically synthesized RNA or DNA.

- The term "encoding" is intended to mean that the subject nucleic acid may be transcribed and translated into the subject protein in a cell, e.g., when the subject nucleic acid is linked to appropriate control sequences such as promoter and enhancer elements in a suitable vector (e.g., an expression vector) and when the
15 vector is introduced into a cell.

The term "polypeptide" is used to mean three or more amino acids linked in a serial array.

- The term "antisense DNA" is used to mean a gene sequence DNA that has a nucleotide sequence homologous with the "sense strand" of a gene when read in a
20 reversed orientation, i.e., DNA read into RNA in a 3' to 5' rather than 5' to 3' direction. The term "antisense RNA" is used to mean a RNA nucleotide sequence (e.g., encoded by an antisense DNA or synthesized complementary with said antisense DNA). Antisense RNA is capable of hybridizing under stringent conditions with an antisense DNA. The antisense RNA of the invention is useful for inhibiting expression of a
25 "target gene" either at the transcriptional or translational level. For example, transcription of the subject nucleic acids may produce antisense transcripts that are capable of inhibiting transcription by inhibiting initiation of transcription or by competing for limiting transcription factors; or, the antisense transcripts may inhibit transport of the "target RNA"; or, the antisense transcripts may inhibit translation of
30 "target RNA".

- The term "sense strand" is used to mean the single stranded DNA molecule from a genomic DNA that is transcribable and translatable into the polypeptide product of the gene. The term "antisense strand" is used to mean the single strand DNA molecule of a genomic DNA that is complementary with the sense strand of the
35 gene.

- 9 -

The term "capable of hybridizing under stringent conditions" is used to mean annealing a first nucleic acid to a second nucleic acid under stringent conditions (defined below). For example, the first nucleic acid may be a test sample, and the second nucleic acid may be the sense or antisense strand of a prosystemin gene.

- 5 Hybridization of the first and second nucleic acids is conducted under stringent conditions, e.g., high temperature and/or low salt content, which tend to disfavor hybridization of dissimilar nucleotide sequences. A suitable protocol involving hybridization in 6 X SSC, at 42°C in aqueous solution followed by washing with 1 X SSC, at 55°C in aqueous solution is provided in the illustrative examples below.
- 10 (Other experimental conditions for controlling stringency are described in Maniatis, T., et al., *Molecular Cloning; A Laboratory Manual*, Cold Springs Harbor Laboratory, Cold Springs, NY, 1982, at pages 387-389; and also in Sambrook, Fritsch, and Maniatis, *Molecular Cloning; A Laboratory Manual, Second Edition*, Volume 2, Cold Springs Harbor Laboratory, Cold Springs, NY, 1989, pages 8.46-8.47.)

- 15 The term "fragment" when used herein with reference to nucleic acid (e.g., cDNA, genomic DNA, i.e., gDNA) is used to mean a portion of the subject nucleic acid such as constructed artificially (e.g., through chemical synthesis) or by cleaving a natural product into a multiplicity of pieces (e.g., with a nuclease or endonuclease to obtain restriction fragments).

- 20 The term "synthetic oligonucleotide" refers to an artificial nucleic acid (e.g., a chemically synthesized nucleic acid) having 9 or more nucleotides.

- The term "systemin polypeptide" is used to mean a polypeptide having an amino acid sequence $R_1R_1QR_1R_2PPR_1R_2R_2R_1PPR_2R_1QR_1R_1$, wherein R_1 is any amino acid, R_2 is lysine or arginine (or derivative thereof), Q is glutamine (or derivative thereof), and P is proline (or any other derivative thereof), e.g., the systemin polypeptide of FIGURE 3; namely, $NH_3-AVQSKPPSKRDPPKMQTD-COO-$. Skilled artisans will recognize that through the process of mutation and/or evolution, polypeptides of different lengths, e.g., with insertions, substitutions, deletions, and the like, may have arisen that are related to the systemin polypeptide of the invention by
- 25 virtue of: a) amino acid and/or nucleotide sequence homology; b) a defensive function in regulating gene expression in response to predators, pathogens, and mechanical injury; and/or, c) the organization of the genomic DNA, as described in Example 6, below. Representative examples of systemin family members in tomato and potato are provided in Example 6-9 (below), and illustrative methods for
- 30 identification of systemin family members in other species, genera, and families of
- 35

- 10 -

plants are also provided in Examples 6-9 (below).

The term "systemin nucleic acid" is used herein to refer to that subset of nucleic acids capable of encoding a systemin polypeptide.

The term "prosystemin polypeptide" is used to mean a precursor polypeptide
5 capable of giving rise to a systemin polypeptide. A representative example is provided
by the prosystemin polypeptide encoded by the cDNA of FIGURE 7 or the coding
region of the genomic DNA of FIGURES 8A, 8B, and 8C. Prosystemin polypeptide is
capable of being cleaved (e.g., chemically or enzymatically) to give rise to systemin.
A representative method for identifying prosystemin genes in different species of
10 plants is provided in Example 9, below.

The term "prosystemin nucleic acid" is used herein to refer to that subset of nucleic acids capable of encoding a prosystemin polypeptide.

Embodiments of the invention described and illustrated below provide systemin
and prosystemin polypeptides, nucleic acids encoding systemin and prosystemin
15 mRNA, cDNA, and genomic DNA, and, including 5' regulatory sequences controlling
transcription of prosystemin gDNA into mRNA. The subject nucleic acids of the
invention are capable of encoding prosystemins that are constitutively synthesized at
a low level and wound-inducible to a high level (see illustrative Example 7, below).
Would-inducible and constitutive low-level expression is provided by regulatory
20 elements within 3000 bp of the 5' region of the systemin gene sequence, the first 104
nucleotides of which are shown in FIGURE 8A. Promoter, enhancer, and other
regulatory elements within the 3000 bp 5' region are useful for insertion into
recombinant plasmids and vectors for controlling gene expression in plants, (i.e.,
genes other than prosystemin). Representative examples of genes that may be linked
25 to the 5' regulatory elements of prosystemin include; genes encoding storage or
nutritionally important proteins, such as vegetative storage proteins, seed storage
proteins, tuber storage proteins and the like; and, genes encoding other plant defense
genes, i.e., other proteinase inhibitors Bt toxin, and the like; genes encoding
regulatory enzymes for metabolic and defensive processes, including phenylalanine
30 amines, HMG CoA reductase and the like; genes encoding commercially important
enzymes in plant suspension culture, such as proteinases, lipases, and the like; and,
genes that regulate flower color.

Purification and physical properties of a representative systemin polypeptide
are disclosed (Example 1, below). Skilled artisans will recognize that the relatively
35 high proportion of hydrophilic amino acids in the prosystemin polypeptide suggest a

- 11 -

variety of conventional approaches to purification that may be used to purify a natural, recombinant, or synthetic prosystemin polypeptide, (e.g., ion exchange chromatography, affinity chromatography, specific ion precipitation, and the like).

The subject amino acid sequence of prosystemin disclosed herein provides amino acid sequences that may be used to construct synthetic peptides of prosystemin or systemin; or, alternatively they may be used to instruct sites at which cleavage of a prosystemin polypeptide will liberate a systemin (e.g., enzymatic cleavage sites in a natural prosystemin or a chimeric recombinant prosystemin protein. In the latter case a chimeric recombinant prosystemin polypeptide may be produced in an expression system, the chimeric protein purified, and then systemin liberated from the chimeric protein by enzymatic cleavage.). Cleaving a prosystemin polypeptide at boundary amino acids produces systemin, e.g., cleaving the prosystemin of FIGURE 7 at both Leu₁₇₈-Ala₁₇₉, (e.g., cleaving with an Leu-Ala-specific endopeptidase; abbreviated, LA peptidase) and at Asp₁₉₆-Asn₁₉₇ (e.g., cleaving with an Asp-Asn-specific endopeptidase; abbreviated, DN peptidase). As an alternative to the LA peptidase, a prosystemin polypeptide may also be cleaved by suitable enzymes at other upstream sites such as Arg₁₇₅-Glu₁₇₆ or Glu₁₇₆-Asp₁₇₇; followed by sequential cleavage of the product with an N-terminal peptidase, i.e., until the LA residues are reached and cleaved. In a similar manner and as an alternative to a DN peptidase, a carboxypeptidase or carboxydipeptidase may be used to sequentially remote amino acids, from the carboxy-terminus until the DN residues are reached and cleaved. Those skilled in the art will recognize that a suitable LA-specific peptidase(s) may be isolated from plant tissues, e.g., by using natural (or synthetic) polypeptide substrates having the prosystemin-systemin boundary amino acid sequences (e.g., L-A and D-N) and assaying for the production of systemin biological activity. In one such illustrative example, a recombinant prosystemin chimeric protein may be synthesized by an expression system and used as a substrate in enzymatic assays to identify and isolate the LA and/or DN peptidase(s).

Those skilled in the art will recognize that the subject prosystemin amino acid sequence may be used for constructing proteinase inhibitors specific for the LA and/or DN peptidases, and such inhibitors may be useful for inhibiting systemin production from prosystemin; thereby inhibiting systemin activation of defense protein production in plants. Skilled artisans will also recognize that LA and DN peptidase may be selected with enhanced ability to liberate systemin from prosystemin (e.g. LA and DN enzymes having increased turnover number, decreased Km, increased Vmax, or

- 12 -

decreased sensitivity to feedback inhibition, and the like). Strains of plants may either be selected, or constructed (*i.e.*, as transgenic plants), having increased LA and/or DN peptidase activity. The subject plants may exhibit increased resistance to predators.

The subject systemin polypeptides of the present invention may also be used
5 for identifying and isolating systemin receptors from plant cells. Those skilled in the art will recognize that the subject polypeptides can be labeled (*e.g.*, with a radioactive label) and conjugated to a photochemical crosslinking agent. The subject conjugated and radiolabeled polypeptides bind to the cellular systemin receptor and photochemical activation forms covalent bonds between the polypeptide and its
10 receptor. When the receptor-polypeptide complex is extracted from the cell it may be isolated and identified by virtue of its label, *e.g.*, the molecular size may be conveniently determined by SDS-PAGE and autoradiography. The subject polypeptides of the invention may also be useful in ligand affinity chromatography for isolating systemin receptors.

15 Embodiments of the invention provide processes for enhancing or inhibiting synthesis of a defense protein in a plant by introducing the subject nucleic acids of the invention into a plant cell. In one representative example enhanced defense protein production may be achieved by inserting prosystemin (or systemin) nucleic acid in a vector downstream from a promoter sequence capable of driving constitutive
20 high-level expression in a plant cell. In the latter case, when the subject vector is introduced into a plant cell the cells containing one or more copies of the subject nucleic acid may exhibit increased synthesis of systemin. When grown into plants the transgenic plants may exhibit increased synthesis of defense proteins, and increased resistance to herbivores, as discussed in more detail in Example 11, below.

25 In another embodiment, the invention provides processes for inhibiting synthesis of defense proteins in a plant by inserting prosystemin antisense nucleic acid in a vector downstream from a promoter sequence. When the latter construct is introduced into plant cells the cells containing one or more copies of the subject nucleic acid may exhibit decreased synthesis of defense proteins. A representative
30 example of a prosystemin antisense vector, and process for inhibiting synthesis of defense proteins is provided in Example 10, below.

Transgenic plants containing the subject antisense nucleic acids of the invention are useful for: a) identifying other mediators that may be present in the prosystemin molecule, (*e.g.*, other mediators that may induce expression of defense
35 proteins or differentiation); b) establishing the extent to which any specific insect

- 13 -

and/or pathogen is responsible for damage of a particular plant. In the latter case the transgenic plants of the invention are useful for assessing the importance of systemin defense mechanisms in production of a plant as a crop.

In still other embodiments, the invention provides transgenic plants constructed
5 by introducing a subject nucleic acid of the invention into a plant cell, and growing the cell into a callus and then into a plant; or, alternatively by breeding a transgenic plant from the subject process with a second plant to form an F1 or higher hybrid (*i.e.*, F2). The subject transgenic plants and progeny may be used to find those plants that contain extra copies of the subject nucleic acid of the invention, and increased
10 expression of prosystemin or systemin. A representative example of a process for producing such a transgenic plant, and breeding it to obtain F1 offspring is provided below in Example 10.

Those skilled in the art will recognize the agricultural advantages inherent in plants constructed to have either increased or decreased expression of systemin
15 polypeptide. For example, such plants may have increased resistance to attack by predators, insects, pathogens, microorganisms, herbivores, mechanical damage and the like. Skilled artisans will also recognize that chemical agents may be developed that will mimic or induce systemin activity (*e.g.*, in a manner similar to methyl jasmonate induction of systemin activity), and these chemical agents may be useful
20 when sprayed on plants in maximizing crop resistance to herbivores, pathogens, and mechanical damage. Representative examples of plants in which the process may be useful include (but are not limited to) tomato, potato, tobacco, corn, wheat, rice, cotton, soybean, alfalfa, rape, poplar trees, pine and fir trees and the like.

The subject nucleic acids of the invention are also useful as oligonucleotide
25 probes (*e.g.*, ³²P-labeled synthetic oligonucleotides constructed to complement portions of prosystemin nucleotide sequence), and restriction fragment probes (*e.g.*, end-labeled restriction fragments of prosystemin cDNA), in Northern and Southern blots for selecting and screening among plants to find natural and mutant strains with increased prosystemin expression and/or genomic copy number. This screening
30 procedure can be useful for identifying plant strains with increased resistance to attack by predators, herbivores, insects, bacteria, fungi, viruses, mechanical damage, and the like.

The subject polypeptides of the invention are useful for inducing monoclonal and polyclonal antibodies that may be used in immunoassays to detect the presence
35 or amount of a prosystemin or systemin polypeptide in plant tissues, extracts, and

- 14 -

fluids (e.g. see E. Harlow and D. Lane, *Antibodies: A Laboratory Manual*, Cold Springs Harbor Laboratory, Cold Springs, N.Y. 1988). The latter immunoassays may prove useful for identifying natural and mutant strains of plants with increased levels of prosystemin or systemin. Strains exhibiting increased levels of the subject

5 polypeptides may have increased resistance to attack by herbivores, *i.e.*, insects, bacteria, fungi, and viruses.

Systemin is a primary polypeptide signal mediating wound-inducible expression of defense genes in distal leaves and, therefore, is the first example of a peptide hormone found in plants. The expression of a recombinant antisense genetic

10 expression construct (*i.e.*, containing an antisense prosystemin cDNA) resulted in an almost complete suppression of systemic wound induced defense protein synthesis in plants. The latter finding provides evidence that systemin is an integral component of the systemic signal transduction system in plants that induces defense protein synthesis in response to attack by predators and the like. It is considered most likely

15 that systemin is the first member to be identified in a systemin family of plant polypeptide hormones. It is considered highly likely that members of the systemin family may regulate developmental events in the meristems, flower tissues, and fruit of plants, *e.g.*, tomatoes and potatoes. Other members of the systemin family may be identified by virtue of their amino acid or nucleotide sequence homology with

20 prosystemin or systemin, or by their ability to hybridize with the subject prosystemin or systemin nucleic acids of the invention. (In this regard, the nucleotide sequences of the exons identified in Example 6, below, may prove useful as oligonucleotide probes for identifying other systemin family members.) In this case the ability of a DNA or RNA to hybridize with the nucleic acid of the invention under conditions of

25 reduced stringency, (*e.g.*, a suitable protocol involving hybridization in 6 X SSC, at 42°C in aqueous solution followed by washing with 1 X SSC, at 55°C in aqueous solution) will be considered a preliminary indication that the DNA or RNA contains a systemin family member. The DNA or RNA may then be sequenced and the sequence compared with the sequence of systemin and prosystemin. (Experimental conditions

30 for controlling stringency are also described in Maniatis, T., et al., *Molecular Cloning; A Laboratory Manual*, Cold Springs Harbor Laboratory, Cold Springs, NY, 1982, at pages 387-389; and, also in Sambrook, Fritsch, and Maniatis, *Molecular Cloning; A Laboratory Manual, Second Edition*, Volume 2, Cold Springs Harbor Laboratory, Cold Springs, NY, 1989, pages 8.46-8.47.) Systemin family members may be recognized

35 by virtue of about 50% to about 100%, or more preferably about 70% to about 100%,

- 15 -

and most preferably about 80% to about 100% homology at the amino acid or nucleotide level, *i.e.*, over a stretch of about 5 or more amino acids or about 15 or more nucleotides.

The foregoing may be appreciated more fully by reference to the following
5 representative examples of the subject compositions and methods provided by the invention.

EXAMPLE 1

Isolation and sequencing of the systemin polypeptide.

Oligogalacturonides were initially considered to be primary candidates as
10 systemic signals for the wound response because they elicit synthesis of antibiotic phytoalexins in plant cells near the sites of infections (10, 11). Oligogalacturonides are released by pectin-degrading that are not found in tomato leaves. In addition, when labeled α -1, 4-oligogalacturonides were applied to wound sites on tomato plants, they were not found to be mobile (E.A. -H. Baydoun and S. Fry, *Planta* 165,
15 269 1985). Thus, oligogalacturonides are probably not involved as systemic mediators of signal transduction in plants, at least with respect to induction of proteinase inhibitor genes in response to wounding.

A search was initiated for systemic signals inducing proteinase Inhibitor I and II genes in tomato leaf extracts. This search led us to identify a polypeptide in tomato
20 leaves that is free of carbohydrates and induces proteinase inhibitor activity when supplied to young tomato plants. The polypeptide was purified using high-performance liquid chromatography (HPLC, see Materials and Methods, below). Inducing activity of the polypeptide was assayed by cutting the petioles of young plants and introducing eluted fractions from column separations into the cut over a
25 period of 30 min. The plants were subsequently transferred to small vials of water, incubated under constant light for 24 hours as described (C.A. Ryan, *Plant Physiol.* 54, 328, 1974), and the amount of proteinase Inhibitor I and II in the leaf juice was quantified by radial immunodiffusion in agar gels that contained rabbit antiserum to Inhibitor I or Inhibitor II (C.A. Ryan, *Anal. Biochem.* 19, 434, 1967; R. Trautman, K.
30 Cowan, G. Wagner, *Immunochemistry* 8, 901, 1971). Over 30,000 young tomato plants were assayed over a 2.5 year period. With the use of this protocol, slightly more than 1 μ g of an active factor (*i.e.*, systemin) was isolated from approximately 60 pounds of tomato leaves.

The elution profile of the preliminary extract of tomato leaves (FIGURE 1) was
35 complex. Several fractions exhibited proteinase inhibitor inducing activity but one

- 16 -

peak (FIGURE 1) was selected for further purification because it contained the highest activity and the best yield from the purification.

After several additional purification steps (see Materials and Methods, below), a major peak that possessed high specific activity was eluted from a strong cation exchange (SCX) HPLC column (FIGURE 2). The properties of the eluted material resembled those of a polypeptide, that is, absorbance in the spectral region appropriate for peptide bonds, total loss of activity and recovery of free amino acids after acid hydrolysis, partial loss of activity in the presence of trypsin and other proteolytic enzymes, and a positive assay result with bicinchoninic acid (P.K. Smith et al., *Anal. Biochem.* 150, 76, 1985). Total amino acid analysis of the bioactive peak eluted from the SCX-HPLC (step 5, Materials and Methods, below) was determined as described below. The amino acid sequence analysis of the active component (conducted as described below; see, Materials and Methods) identified its length and determined the sequence of FIGURE 3: NH₃-AVQSKPPSKRDPPKMQTQTD-COO⁻). No significant similarities were found to known protein sequences and the polypeptide was named "systemin" (Protein Identification Resource release 26; Pearson/Lipman FASTA program at the Molecular Biology Computer Research Resource, Harvard Medical School). The sequence is a palindrome: xxQxBPPxBBxPPBxQxx (x, any amino acid residue; B, Lys or Arg; Q, Gln; P, Pro). A synthesized polypeptide of identical sequence to the systemin sequence (prepared as described below; see, Materials and Methods) eluted from the C18 (step 2) column with the same retention time as the native polypeptide.

Materials and Methods:

Purification and isolation of the polypeptide inducer of defense proteins:

Step 1: Approximately 2 kg of tomato leaves *Lycopersicon esculentum* (v. Castlemart) were harvested from 20-day-old plants, grown under cycles of 17 hours light at 28°C and 7 hours dark at 18°C. Leaves were homogenized in a Waring blender for 5 min with distilled water (total volume of 4 liters) and filtered through four layers of cheesecloth. The liquid was adjusted to pH 4.5 with HCl and centrifuged at 1000g for 10 min. The supernatant was adjusted to pH 6.1 with 10N NaOH, centrifuged at 10,000g for 10 min at 20°C, and decanted through Whatman #4 filter paper. The filtrate was chromatographed on DEAE cellulose, followed by reversed-phase C18 flash chromatography, Sephadex G25 gel filtration, and then CM Sephadex chromatography.

The DEAE cellulose column (Whatman DE52, 5.9 cm by 15 cm) was

- 17 -

equilibrated in 1M ammonium bicarbonate and washed exhaustively with distilled water. The material eluting in the void volume was collected and stored overnight at 4°C. TFA was added dropwise to the stored elute to a final concentration of 0.2% (v/v); the solution was then clarified by centrifugation at 20,000g for 5 min at room temperature. The supernatant was loaded onto a reversed-phase flash column (C18, 40µm, 3 cm by 25 cm) previously equilibrated with aqueous 0.1% TFA. The column was eluted with the use of compressed nitrogen at 8 psi. After the sample was loaded, the column was washed with 200 ml 0.1% TFA; the retained material was then eluted with successive washes of 20, 40, and 60% methanol in 0.1% TFA. The methanol was removed with a rotary evaporator and the remaining liquid was frozen and lyophilized. Two kilograms of leaf material yielded about 1 g of crude material containing systemin. The procedure was repeated 15 times. Samples (approximately 4 g) of crude material dissolved in 20 ml water and adjusted to pH 7.8 with 10M ammonium hydroxide were loaded onto a G25 Sephadex column (4 cm by 44 cm) that was equilibrated with 50 mM ammonium bicarbonate, pH 7.8. The material eluting at and just after the void volume was recovered and lyophilized. Four identical runs through the entire procedure produced 1.25 g of partially purified systemin. The 1.25 g was dissolved in 500 ml H₂O, the pH was adjusted to 6 with 1M NaOH, and the sample was applied to a CM Sephadex column (2 cm by 17 cm) and washed with 0.01 M potassium phosphate, pH 6. The activity was retained by the CM Sephadex, eluted with 250 mM ammonium bicarbonate, and lyophilized. The total yield of proteins in this step was 190 mg.

Step 2: The active fraction (190 mg) recovered from step 1 was dissolved in 10 ml 0.1% TFA, centrifuged at 20,000g for 5 min, filtered, and chromatographed on a reversed-phase C18 column. Five microliters of each 2-ml eluted fraction was diluted to 360 µl with 154 mM sodium phosphate, pH 6.5, and assayed for proteinase Inhibitor I inducing activity (x in Figure 1). Four plants were assayed per fraction. The material was injected into a semi-preparative reversed-phase C18 column (Vydac, Hesperia, CA, Column 218 TP510, 10mm by 250mm, 5-µm beads, 300A pores). Solvent A consisted of 0.1% TFA in water. Solvent B consisted of 0.1% TFA in acetonitrile. Samples were injected in solvent A and, after 2 min, a 90 minute gradient to 30% solvent B was begun for elution. The flow rate was 2 ml/min and eluted peaks were monitored at 225 nm. Several peaks of biological activity were found (as described below). The major peak of activity (shown in black in Figure 1) resided in tubes 43 to 46, which were pooled and lyophilized. Total protein content of the pooled fractions

- 18 -

was estimated at 2.5 mg.

Step 3: The total material recovered in step 2, above (2.5 mg), was subjected to strong cation exchange HPLC on a poly-SULFO-ETHYL Aspartamide (SCX) column (4.6 mm by 200 mm, 5 μ m, The Nest Group, Southborough, MA) with the use of the following solvent systems: Solvent A, 5mM potassium phosphate, pH 3, in 25% acetonitrile; solvent B, 5mM potassium phosphate, 500 mM potassium chloride in 25% acetonitrile, pH 3. The sample was dissolved in 2 ml of solvent A, filtered, and applied to the column. After a 5-minute wash with solvent A, a 60-min gradient to 50% B was applied. The flow rate was 1 ml/min, and the elution profile was monitored by absorbance at 210 nm. The active fractions, tubes 35 to 38, were pooled and reduced in volume to 1 ml by vacuum centrifugation.

Step 4: The pooled fractions from step 3 were subjected to reverse-phase C18 HPLC in 10mM potassium phosphate, pH 6. Chromatography was performed on a Beckman Ultrasphere Ion pair column (4.6 mm by 250 mm, C18, 5 μ m). Solvent A was 10 mM potassium phosphate, pH 6, and solvent B was 10 mM potassium phosphate, pH 6, containing 50% acetonitrile. The active fractions, tubes 39 to 42, were pooled and vacuum centrifuged to a final volume of 1 ml. This fraction was applied to the same column as the previous run but under the solvent and gradient conditions of step 2. The sample was adjusted to pH 3 with TFA, filtered through a 0.45- μ m syringe filter and chromatographed at a flow rate of 1 ml/min. The peaks of protein were detected at 212 nm. The fractions containing activity, eluting at 53.5 to 56.5 min, were pooled and vacuum centrifuged to a volume of 1 ml.

Step 5: The active fraction from step 4 was subjected to SCX-HPLC with the same column and conditions as used in step 3, except that the gradient was shallower, i.e., the column was run at 0% B for 5 min at which time a gradient to 30% B in 120 min was started. Fractions (0.5 ml) were diluted as in FIGURE 1 and assayed for proteinase Inhibitor I inducing activity (x in Figure 2). The biologically active fractions of the system peak (shown in black in Figure 2) were collected and analyzed for amino acid content and sequence. The profile was detected by absorbance at 210 nm. Fractions eluting at 76 to 78.5 min were pooled and vacuum centrifuged to reduce the volume to 1 ml.

Step 6: The step 5 fraction was desalted on a C18 HPLC column under the conditions of step 2. A 60-minute gradient to 30% solvent B was employed. The fractions containing the activity peak eluted at 55.0 to 58.0 min and were pooled and concentrated by vacuum centrifugation to 0.5 ml. The sample contained

- 19 -

approximately 1 μ g of protein, as estimated by amino acid content after acid hydrolysis. The biological activity of the sample had the potential to induce maximal accumulation of proteinase inhibitors in 40,000 tomato plants, (*i.e.*, approximately 40,000-fold purified). This sample was used for amino acid analysis and sequence
5 determination.

Amino acid analysis:

The bioactive peak eluted from the SCX-HPLC column (*step 6*) was dried in 6 by 50 mm glass tubes and hydrolyzed in the HCl vapor. The hydrolysates were derivatized with phenylisothiocyanate and analyzed by reverse-phase chromatography
10 on 30 cm by 0.39 cm columns (Picotag, Millipore) according to the manufacturer's suggestions.

Amino acid sequence analysis:

The amino acid sequence of the bioactive peak-eluted from the SCX-HPLC column (*step 6*) was determined by established methods (D.J. Strydom et al.,
15 *Biochemistry* 25, 945, 1985; B.A. Bidlingmeyer, S.A. Cohen, T.L. Tarvin, *J. Chromatogr.* 336, 93, 1984). Briefly, sequencing was performed on a Beckman model 890 spinning cup instrument, equipped for microsequencing, as recommended by the manufacturer, except that 0.1% water was added to the anhydrous heptafluorobutyric acid (HFBA) and 0.1% ethanethiol was added to the 25% trifluoroacetic acid (TFA). Identification of
20 the phenylthiohydantoin amino acids was by reverse-phase chromatography on an octadecylsilane column (IBM, Inc.; 30 cm by 0.46 cm, 3- μ m particle size) with the use of the gradient system described (D.J. Strydom et al., *supra*). Abbreviations for the amino acid residues are: A, Ala; C, Cys; D, Asp; E, Glu; F, Phe; G, Gly; H, His; I, Ile; K, Lys; L, Leu; M, Met; N, Asn; P, Pro; Q, Gln; R, Arg; S, Ser; T, Thr; V, Val; W, Trp;
25 and Y, Tyr.

Synthesis of a synthetic systemin polypeptide:

A polypeptide corresponding to residues 2 through 18 (FIGURE 3) was synthesized with the use of 9-fluorenylmethyl chloroformate (F-moc) solid phase chemistry with an Applied Biosystems Inc. Model 431A synthesizer on a p-methyl
30 benzyhyrlyamine resin according to the manufacturer's protocol. The F-moc derivative of [14 C] Ala (New England Nuclear) was synthesized (J. Stewart and J. Young, *Solid Phase Peptide Synthesis* (Pierce Chemical Co., Rockford, IL, ed. 2, 1984), pp.67-68) and added to the NH₂- terminal residue of the polypeptide; the [14 C] Ala-polypeptide was then cleaved from the resin. The radioactive peptide was purified by C18 HPLC.
35 The specific activity of the synthetic systemin polypeptide was 19.0 μ Ci/ μ mol.

- 20 -

EXAMPLE 2

Systemic translocation of radiolabeled systemin polypeptide.

The synthetic systemin polypeptide (described above) was tested for bioactivity and was found to be as effective as the native systemin polypeptide (purified above to step 6) for inducing the synthesis and accumulation of both Inhibitor I and II proteins when supplied to the cut stems of young tomato plants (FIGURE 4). About 40 fmol of the polypeptide per plant was required to produce half maximal accumulation of Inhibitors I and II, which represents about 10^5 times more activity on a molar basis than the previously reported PIIF oligogalacturonide inducers derived from plant cell walls. The coordinate induction of synthesis of Inhibitor I and Inhibitor II proteins in response to the systemin synthetic polypeptide (FIGURE 4) is similar to a normal plant wound response that is transcriptionally regulated. This suggests that the polypeptide is activating the same signal transduction pathway activated by wounding (T. Green and C.A. Ryan, 1972, *supra*; J.S. Graham et al., 1986, *supra*), by oligosaccharides (P. Bishop et al., 1981, *supra*; M. Walker-Simmons et al., 1983, *supra*), and by methyl jasmonate (E.E. Farmer et al., 1990, *supra*).

The systemin polypeptide, unlike the prior reported oligogalacturonides, is transported out of wounds to distal tissues. ^{14}C -labeled polypeptide was synthesized (as described above, Example 1) and placed on fresh wounds of tomato plants. Within 30 min the radioactivity had moved throughout the leaf, and within 1 to 2 hours radiolabeled systemin was identified by HPLC (FIGURE 5) in the phloem exudate (expressed from the plant according to the method described by R.W. King and J.A. ZeeVaart, *Plant Physiol.* 53, 96, 1974). The polypeptide was named "systemin" because of its mobility through phloem.

As well as being inducible by wounding in leaves, the genes for proteinase inhibitors I and II are developmentally regulated in the meristems, flower tissues, and fruit of tomato species, and in potato tubers. Thus, it is likely that these developmental events may be mediated by systemin or by similar polypeptides that are members of the systemin family.

FIGURE 4 shows the results of experiments in which systemin synthetic polypeptides induced synthesis and accumulation of proteinase Inhibitor I (●) and II (○) in the excised leaves of young tomato plants. The leaves were incubated in solutions containing the synthetic systemin polypeptide and the proteinase inhibitors were assayed as described above in Example 1. Each data point was obtained from assays of the leaves of 36 tomato plants.

- 21 -

EXAMPLE 3*Molecular cloning prosystemin.*

A prosystemin cDNA was isolated by screening a primary cDNA library synthesized from tomato leaf mRNA as follows:

5 Poly A + mRNA was purified from tomato leaves using oligo dT columns (Pharmacia). cDNA was synthesized using the Stratagene cDNA synthesis system and was cloned into lambda ZAP vector arms (Stratagene). Approximately 80,000 primary library recombinants were screened, by duplicate plaque lifts, using a degenerate oligonucleotide probe, termed "SP1", based on the amino acid sequence
10 of the carboxy terminus of systemin (*i.e.* PPKMQTN; amino acids 190 to 196, as numbered in FIGURE 6, excluding the last nucleotide residue of the Asp₁₉₆ codon). The hybridization conditions for screening were: 6X SSC; 1X Denhardt's solution; 100 µg/ml, yeast tRNA; 0.5% sodium pyrophosphate and approximately 4x10⁷ cpm of ³²P-end-labeled SP1. Hybridization was at 37°C for 36 hr. The filters were washed in
15 several changes of 5X SSC, 0.1% SDS at 48°C for one hour. Approximately 50 positive clones were identified and rescreened using a second degenerate oligonucleotide, termed "SP2", corresponding to the amino terminus of systemin (AVQSKP; amino acids 179 to 184, as numbered in FIGURE 6, excluding the last of the Pro₁₈₄ codon). The hybridization and wash conditions were identical to those
20 used for SP1 except that the wash temperature for SP2 was 40°C. Of the initial positive clones only one hybridized to the SP2 probe. Fragments of the prosystemin cDNA, termed "pSYS 1", were subcloned into Bluescript® plasmids, single-stranded DNA was rescued and sequenced on both strands by dideoxy sequencing using Sequenase (USB; Sanger et al. *PNAS* 73, 5463, 1977). Sequencing of the SP2
25 positive clone established that it encoded the systemin polypeptide within the larger protein which was called "prosystemin". The prosystemin cDNA was not full-length, beginning at nucleotide 112 as numbered in FIGURE 6.

The prosystemin cDNA consisted of 839bp with an open reading frame encoding 197 amino acids. The reading frame remained open to the 5'-end of the
30 clone, and since Northern blot analysis indicated that the systemin mRNA was 1Kb in size, we concluded that the cDNA was missing approximately 100bp at the 5'-end. The complete prosystemin mRNA sequence was subsequently determined by sequencing the prosystemin gene (as described below in Example 6) and mapping the transcriptional start site (FIGURE 7; Example 6). The experiments described in
35 Example 6, below, established the length of the open reading frame as 600 base pairs

- 22 -

encoding a prosystemin protein of 200 amino acids. The identification of the initiating methionine codon was made on the basis of two criteria; the presence of multiple stop codons immediately 5' to the methionine codon and the presence of an adjacent sequence similar to the plant consensus sequence for translational initiation (H.A. Lutcke et al., 1987, *EMBO Journal* 6:43).

EXAMPLE 4

Structure and properties of prosystemin.

Based on the cDNA sequence (FIGURE 7), systemin is located close to the carboxy terminus of the 23 kDa prosystemin protein (FIGURE 7, amino acid residues 179 to 196; corresponding to nucleotides 639 to 699). The amino acid composition of prosystemin is unusual in that it contains a high percentage of charged amino acids; aspartic acid (10%), glutamic acid (17%), lysine (15%), but very few hydrophobic amino acids. In consequence, prosystemin is a markedly hydrophilic molecule. Analysis of the prosystemin sequence failed to reveal a hydrophobic region at the amino terminus that resembles a leader peptide. The post-translational processing pathway and site of sub-cellular compartmentalization of prosystemin remain to be determined. A search of the EMBL and GeneBank data base, with both the cDNA and deduced protein precursor sequences failed to reveal significant homology to any of the listed sequences.

Although the 18-amino acid systemin sequence occurs only once within the precursor, close to the carboxy terminus, other sequence elements are repeated. There is a short (6 to 9 amino acids), imperfect repeat occurring five times within the prosystemin sequence (crosshatched underlining, FIGURE 7). This observation suggests that at least part of the prosystemin gene may have evolved by multiple gene duplication/elongation events, a conclusion which is supported by the structure of the gene.

EXAMPLE 5

Proteolytic processing sites in prosystemin.

The putative processing sites bordering systemin are shown in FIGURE 7 (*i.e.* amino acid residues 178 and 197). The Leu (178) and Asn (197) processing sites do not conform to the consensus sequence for the endoproteolytic processing sites flanking bioactive peptides within animal prohormone precursors (*e.g.*, see animal sites in R.B. Harris, 1989, *Arch. Biochem. Biophys.* 275(2):315 (1989). The minimum animal consensus sequence consists of a pair of basic amino acids which immediately precede the site of cleavage. In addition, the dibasic pair is often preceded, at a

- 23 -

distance of two or three amino acids, by a single basic amino acid. The animal consensus sequence is, however, found once within the prosystemin sequence, at residues 183-188 (LysProProSerLysArg, FIGURE 7), which is a part of the mature systemin polypeptide. It is conceivable that the half-life of systemin is regulated by further processing at this site, e.g., to yield an 8 amino acid carboxy-terminal peptide.

In animal systems prohormones are often processed to yield multiple bioactive peptides (J. Douglass, O. Civelli and E. Herbert, 1984, *Ann. Rev. Biochem.* 53:665; L.J. Jung and R.H. Scheffler, 1991, *Science* 251:1330) and members of the systemin family of plant polypeptide hormones may be subject to similar processing mechanisms.

EXAMPLE 6

Structure of the prosystemin gene and systemin gene family.

A primary library of 700,000 recombinants was plated on the bacterial strain (P2) PLK-17 (Stratagene) and screened, by duplicate plaque lifts, with nick-translated prosystemin cDNA. Hybridization was carried out as described below. A single positive clone was identified and purified: The gene was located on a 18Kb genomic DNA fragment from which it was sub-cloned into Bluescript® plasmids. A series of overlapping, deletions spanning most of the gene were produced using the Mung Bean/Exonuclease III system (Stratagene). Each deletion product was cloned into a Bluescript® phagemid from which single-stranded DNA was rescued for use as the sequencing template. The gene was sequenced by the dideoxy method of Sanger (*supra*) using Sequenase (U.S.B.). The sequence was completed using custom-made oligonucleotide primers. The deduced sequence was then confirmed using custom-made oligonucleotide primers (as described in the Materials and Methods below).

The sequence of the prosystemin gene is presented in FIGURES 8A-8C. The gene is composed of 4526 bp comprising a 104 bp 5'-untranslated region, a 4176 bp coding region and a 246 bp 3'-untranslated region. A striking feature of the sequence is that it is 76% A:T-rich. The structure of the prosystemin gene is depicted in FIGURE 9A; Southern blot analysis is shown in FIGURE 9B.

Within the prosystemin gene ten introns interrupt the coding region (FIGURE 10). In consequence, the exons are small, ranging, in size from 34 bp (exon 1) to 90 bp (exon 10). The first 10 exons are organized as five pairs while the sequence encoding systemin is located on the final, unpaired exon. To investigate the relationship between the exon pairs we aligned the sequences of the first exon of each pair (exons 1, 3, 5, 7, 9) and, separately, we aligned the sequences of the second exon of each pair (exons 2, 4, 6, 8, 10), as shown in FIGURES 11A AND 11B.

- 24 -

The sequences within the first exon group are all homologous to each other as are the sequences within the second exon group. There is no significant sequence homology between the two groups. These observations suggest that the five exon pairs arose by successive duplications from a common ancestor. That the individual
5 exons within a pair are not homologous to each other suggests that the ancestral unit from which the gene is derived was a structure corresponding to the exon pair rather than to the individual exons within a pair.

In contrast to the set of five homologous exon pairs, the exon encoding systemin (exon 11) does not show significant sequence homology with any other part
10 of the prosystemin gene. This observation suggests either that the exon encoding systemin may have arisen separately from the rest of the gene, to which it was subsequently added, or that the exon encoding systemin arose from the same ancestral sequence as the other exons but subsequently evolved more rapidly.

Repeated amino acid sequences within prosystemin and the systemin gene family.

15 The amino acid sequence of prosystemin, like that of the gene, is highly repetitive. A short oligopeptide sequence palindrome was identified above in systemin (Example 1) and a similar theme occurring five times within prosystemin that is encoded by the first exon of each of the five homologue pairs. In addition, the presence of three different, tandemly-repeated polypeptide elements within
20 prosystemin provides a clue about the evolution of the prosystemin gene.

The tandemly repeated elements occur once within the amino-terminal half of prosystemin and once within the carboxy-terminal half of prosystemin. The polypeptide elements within the amino-terminal half of the precursor have been labelled Rep A, Rep B and Rep C and their repeats within the carboxy-terminal half of
25 the molecule have been labelled Rep 2A, Rep 2B and Rep 2C. The sequences of these repeated polypeptide elements are shown aligned in FIGURE 12. The locations of the repeats within prosystemin are shown in FIGURE 13A, while the locations within the prosystemin gene of the DNA sequences encoding the polypeptide repeats are shown in FIGURE 13B. As can be seen from this comparison the Rep a, Rep B and
30 Rep C polypeptides are encoded by two exon pairs (namely, three plus four and five plus six) while Rep 2A, Rep 2B and Rep 2C are also encoded by two other exon pairs (namely, seven plus eight and nine plus ten).

The observations thus suggest that a set of polypeptide sequences, encoded by two pairs of exons, has been duplicated as one step in the evolution of the
35 prosystemin gene. It would then appear that the ancestral gene for prosystemin was

- 25 -

subject to a tandem duplication of a structure composed of at least two pairs of exons. This conclusion precludes a model in which the systemin gene evolved simply by successive duplications of a single exon pair.

Intron boundaries may be shifted within repeated DNA sequences.

5 The amino acid sequences between the polypeptide Rep A, Rep B and Rep C regions, that was found in the amino terminal half of prosystemin, were not duplicated in the carboxy terminal half of the molecule where repeats Rep 2A, Rep 2B and Rep 2C are almost contiguous (FIGURE 13A). This observation is largely explained by the fact that exon 7 (encoding element Rep 2A), is truncated at its 3'-end
10 *(i.e., with respect to the sequence of exon 3, encoding element Rep A).* Similarly, exon 9 (encoding element Rep 2C), is truncated at its 5'-end *(i.e., compared to exon 5, encoding element Rep C).*

 In the case of exon 7, the truncation did not arise by deleting a short section of the gene, but by shifting the position of the intron boundary within the existing
15 sequence. Comparing the sequence around the intron junction at the 3'-end of exon 7 with the corresponding junction sequence at the 3'-end of exon 3 (FIGURE 14), it is evident that the sequence corresponding to the 3'-end of the exon 3 constitutes the 5'-end of the intron between exons 7 and 8. This structure may have arisen by either the elongation of exon 3 or the shortening of exon 7. It is not clear if the same
20 process resulted in the truncation of the 5'-end of exon 9.

Materials and Methods:

 Primer extension was performed using a gel-purified oligonucleotide consisting of bases 82 to 111 of the antisense strand of the cDNA sequence shown in FIGURE 7. Total RNA was extracted from young tomato plants 4 hr after wounding and poly
25 A+ mRNA was isolated using oligo dT columns (Pharmacia). Three picomoles of oligonucleotide were end-labelled using γ -ATP at a specific activity of 6000/Ci/mmol. 2×10^6 cpm of labelled oligonucleotide were annealed to 4 μ g polyA+ mRNA by heating to 85°C for 10 min then hybridizing overnight at 30°C in a solution of 40 mM PIPES, pH 6.4; 1 mM EDTA; 0.4 M NaCl; 80% formamide. The annealed nucleic acids
30 were ethanol precipitated and resuspended in 30 μ l of a solution containing: 50 mM Tris, pH 7.5; 75 mM KCl; 10 mM dithiothreitol; 3 mM $MgCl_2$; 500 μ M of each dNTP; 100 μ g/ml bovine serum albumin. Thirty units of M-MLV Reverse Transcriptase (Stratagene) and one unit of RNase Block II (Stratagene) were added and the reaction mixture was incubated for 90 min at 37°C. At the end of the reaction one μ l of 0.5 M
35 EDTA, pH 8.0 and one μ l of DNase-free RNase A (10 mg/ml) were added to the

- 26 -

reaction mixture and incubated for a further 30 min at 37°C. The reaction mixture was phenol extracted, ethanol precipitated and resuspended in four µl TE buffer (10 mM Tris-HCL, pH 7.5, 0.1 mM EDTA, pH 8.0) to which six µl of formamide loading buffer (80% formamide; 10 mM EDTA, pH 8.0; 1 mg/ml xylene cyanol; 1 mg/ml bromophenol blue) were subsequently added. Two µl of the resuspended products were analyzed on a 6% acrylamide/8M urea sequencing gel. The size standards were sequencing products generated using the primer extension oligonucleotide as primer and single-stranded DNA derived from the 5'-end of the prosystemin gene as template. Sequencing was carried out using Sequenase (USB) following the manufacturer's instructions for generating sequencing products close to the primer.

Mung Bean Nuclease analysis was carried out using a 400bp ScaI-NdeI fragment spanning the 5'-end of the prosystemin gene. The NdeI site is located within the first exon of the systemin gene. The NdeI end of the fragment was end-labelled to a specific activity of 6×10^6 cpm/µg and approximately 10^6 cpm were mixed with 4 µg of the same poly A+ RNA stock used in the primer extension experiment. The mixture was desiccated and resuspended in 15 µl of hybridization buffer. The mixture was covered with mineral oil, heated to 82°C for 6 min then hybridized overnight at 37°C. The sample was then mixed with 200 µl of ice-cold Mung Bean Nuclease buffer (30 mM sodium acetate (pH 5.0), 50 mM sodium chloride, 1 mM zinc chloride, 5% (v/v) glycerol) to which 10 units of Mung Bean Nuclease (Stratagene) were added. The mixture was incubated for 30 min at 12°C then extracted with an equal volume of a 1:1 mixture of phenol:chloroform. The digestion products were coprecipitated with 1 µg of yeast tRNA and resuspended in 4 µl TE buffer plus 6 µl formamide loading buffer. Three µl of the resuspended digestion products were analyzed on a 6% acrylamide/8M urea gel. Size markers were generated by using single-stranded DNA corresponding to the 5'-end of the gene as template. The sequencing primer was a 19-mer corresponding to the first 19 bases (antisense strand) at the 3'-end of the ScaI-NdeI probe fragment.

EXAMPLE 7

Wound-inducible expression of the prosystemin gene.

In considering the role of systemin as a mobile signal that activates proteinase inhibitor genes in response to wounding, we investigated the possibility that the prosystemin gene, itself, might be wound-inducible. Northern blot analysis was used to examine the levels of prosystemin mRNA and Inhibitor I mRNA in leaves of unwounded and wounded tomato plants (FIGURE 15A). Thirty-two young tomato

- 27 -

plants were wounded three weeks after germination. The plants had an upper and a lower leaf and a small apical leaf. The lower leaf was wounded and mRNA was isolated from the upper (unwounded) leaf at the following time points after wounding: 0.5, 1.5, 3, 6, 9, 12, and 24 hours. Four plants were used for each time point. Total RNA (5 µg) from each time point was electrophoresed on a 1.4% agarose-formaldehyde gel and blotted onto nitrocellulose. The blot was probed simultaneously with nick-translated prosystemin (SYS) and Inhibitor I (Inh-1) cDNAs (see Materials and Methods, below). Prosystemin mRNA was found to accumulate in both wounded and unwounded leaves of wounded tomato plants, demonstrating that prosystemin mRNA, like Inhibitor I mRNA, is systemically wound-inducible. Prosystemin mRNA reached the highest levels at three to four hours after wounding while Inhibitor I mRNA was most abundant eight to ten hours after wounding. Unlike the proteinase Inhibitor I message, which is absent in the leaves of unwounded tomato plants, a low level of prosystemin mRNA was detected in the leaves of unwounded plants. Low, constitutive expression of the prosystemin gene in leaves may provide a continuous supply of systemin, allowing the plant to immediately respond to wounding.

The wound-induced accumulation of prosystemin mRNA and, presumably, prosystemin and systemin in the unwounded tissue may amplify the ability of the plant to react to subsequent damage. Continued damage by insect attacks would, therefore, liberate more systemin from the newly synthesized precursor than did the initial wounds, resulting in higher levels of proteinase inhibitor synthesis as the attacks persist.

Since the initial rate of accumulation of prosystemin mRNA was faster than that of Inhibitor I mRNA in response to wounding (FIGURE 15A), some aspects of the signal transduction pathways activating the two genes may differ. Additional signals may be responsible for the different rates of accumulation or the signal transduction pathways may respond to the same signals but with different sensitivities.

Materials and Methods:

Nick-translation was performed using the NEN DuPont nick-translation system according to the manufacturer's instructions. Hybridization was carried out under the following conditions: 50% formamide; 5x Denhardt's; 5X SSPE; 0.1% SDS; 100 µg/ml sheared salmon sperm DNA; 1 µg/ml poly A and nick-translated DNA probe of specific activity approximately 10^9 cpm/µg. Unless otherwise stated, blots were washed in 1 X SSC, 0.1% SDS at 65°C.

35

- 28 -

EXAMPLE 8*Distribution of prosystemin mRNA throughout the plant.*

Prosystemin mRNA is found throughout the aerial parts of the plant but not in the roots (FIGURE 15B). Total RNA was extracted from the following parts of an unwounded, fully-grown tomato plant: root (R); stem (St); petiole (Pt), leaf (Le), sepal (Se), petal (Pe), stamen (Sm) and pistil (Pi)(FIGURE 15B). Total RNA (5 μ g) from each sample was electrophoresed and blotted as described in Example 7. The blot was probed with nick-translated prosystemin cDNA (as described in Example 7, above).

The highest constitutive levels of prosystemin mRNA are seen in the flower parts, a feature which is also characteristic of the distribution of Inhibitor I and Inhibitor II mRNAs. The general distribution of prosystemin mRNA (at least in the parts of the plant above ground), is consistent with the proposed role of systemin as a wound signal, since wounding of any aerial part of the plant would be expected to result in the systemic induction of proteinase inhibitor synthesis. The apparent absence of prosystemin mRNA in the roots is surprising since we have observed the induction of proteinase inhibitor synthesis in tomato leaves in response to wounding of the roots. It is possible that roots either contain very low levels of prosystemin mRNA, undetectable in the assay, or they employ a different wound signal(s) (e.g., a different systemin gene family member) to activate proteinase inhibitor genes in the leaves. It is also conceivable that prosystemin is transported from leaves to roots where systemin is released in response to wounding.

EXAMPLE 9*Species distribution of prosystemin gene homologues.*

To determine if prosystemin gene homologues are found in other plant species, Southern and Northern blot analysis was performed on genomic DNA and total RNA from three species known to possess wound-inducible proteinase inhibitors: potato, *Solanum tuberosum*, var. Russett Burbank (C.A. Ryan, 1968, *Plant Physiol.* 43, 1880), tobacco, *Nicotiana tabacum*, var. Xanthi (G. Pearce, results in preparation), and alfalfa, *Medicago sativa*, var. Vernema (W.E. Brown and C.A. Ryan, 1984, *Biochemistry* 23:3418; W.E. Brown, K. Takio, K. Titani, C.A. Ryan, 1985, *Biochemistry* 24:2105); and, as a control, from one species (*Arabidopsis thaliana*, var. Columbia), which is not known to possess wound-inducible proteinase inhibitors.

Southern blot analysis of the species distribution of prosystemin gene homologues is shown in FIGURE 9C. Genomic DNA (5 μ g) from tomato (FIGURE 9C, lane 1), potato (FIGURE 9C, lane 2), tobacco (FIGURE 9C, lane 3), alfalfa (FIGURE 9C,

- 29 -

lane 4), and *Arabidopsis* (FIGURE 9C, lane 5) was digested with EcoR1; restriction fragments were separated by electrophoresis on a 0.8% agarose gel; and, fragments of prosystemin were visualized by blotting to nitrocellulose and probing with nick-translated prosystemin cDNA. The blot was washed at 55°C under moderately stringent conditions. Of the four plant species analyzed, a homologue of the prosystemin gene was identified under moderately stringent conditions only in potato (the nearest relative of the four species to tomato). A potato mRNA species was also identified that hybridized to tomato prosystemin cDNA and which comigrated with the tomato prosystemin mRNA. The nucleotide sequence of the gene in tobacco, alfalfa and *Arabidopsis* may have diverged from that of the tomato gene to the extent that it can no longer be detected by hybridization under stringent conditions with the tomato prosystemin cDNA. This interpretation is favored by the findings that a homologue could not be detected at greatly reduced hybridization and wash stringencies. Analysis of extracts from other plant genera should provide further insights into the distribution and evolution of members of the systemin gene family.

EXAMPLE 10

Antisense suppression of the prosystemin gene.

To determine if the prosystemin gene product has an important role in the systemic signal transduction leading to the expression of proteinase inhibitor genes in tomato leaves, a prosystemin antisense DNA was constructed and was used to transform tomato plants. The chimeric antisense DNA was composed of prosystemin cDNA, in the antisense 3' to 5' orientation, under the control of the constitutive CaMV 35S promoter and inserted into the binary vector pGA643.

Materials and Methods:

Strand-specific, radiolabeled RNA probes were produced from the prosystemin cDNA using T3 and T7 RNA polymerases (Stratagene) according to the manufacturer's instructions.

The antisense DNA construct was transformed into *Agrobacterium* strain LBA4404 and the recombinant bacteria were used to transform tomato var. Better Boy. As controls for the primary transformants, tomato plants were transformed with the binary vector alone. Eighteen antisense plants and twenty one controls were regenerated. Three weeks after the transformed plants had been transferred to soil the lower leaves on each plant were extensively wounded and the levels of wound-inducible proteinase Inhibitors I and II were determined in the expressed juice of upper leaves twenty four hours later (C.A. Ryan, 1967, *Anal. Biochem.* 19:434; R.

- 30 -

Trautman, K.M. Cown, G.G. Wagner, 1971, *Immunochemistry* 8:901). None of the plants were producing either Inhibitor I or Inhibitor II in their leaves prior to wounding. Of the 18 plants containing the antisense gene, 11 plants produced Inhibitor I at less than 40% of the mean control level of 126.7 ± 18.6 $\mu\text{g/ml}$ leaf juice and Inhibitor II at less than 30% of the mean control level of 164.7 ± 18.6 $\mu\text{g/ml}$ leaf juice.

FIGURE 16A shows Northern blot analysis of total RNA isolated from one of the antisense plants, designated 1A4, using both sense and antisense-specific single-stranded RNA probes. Two samples of total RNA (5 μg) were electrophoresed and blotted as described above. The samples were probed separately with radiolabeled RNA probes specific for sense (FIGURE 16A, lane 1) and antisense (FIGURE 16A, lane 2) prosystemin mRNA (see Examples 6-8, above).

In the wounding experiment the distal leaves of plant 1A4 expressed Inhibitor I at 42 $\mu\text{g/ml}$ leaf juice and Inhibitor II at 41 $\mu\text{g/ml}$ leaf juice in response to wounding. The antisense RNA appeared as a band at approximately 1.7 kilobases (FIGURE 16A, lane 2) compared to the prosystemin mRNA at 1 Kb (FIGURE 16A, lane 1). Southern blot analysis showed that plant 1A4 contained a single copy of the antisense construct. This conclusion was confirmed by self-fertilizing plant 1A4 and analyzing 28F1 progeny by Southern blot analysis. Seven (one quarter) of the 28 F1 progeny did not inherit the antisense construct. This experiment also demonstrated that the antisense construct was stably inherited in the F1 generation.

To demonstrate that the antisense phenotype segregated with the antisense construct, the levels of Inhibitors I and II in the distal leaves of the 28 F1 plants were measured before wounding, and 24 hr after wounding. FIGURE 16B graphically depicts wound-induced accumulation of proteinase Inhibitor I and FIGURE 16C depicts wound-induced accumulation of proteinase Inhibitor II in the distal leaves of F1 antisense plants (unshaded bars) and untransformed controls (solid bars). Antisense plant 1A4 was self-fertilized and the amounts of wound-induced proteinase Inhibitors I and II in the distal leaves of three-week old F1 progeny were measured by radial immunodiffusion assay (as described below). The plants had an upper and a lower leaf and a small apical leaf. The lower leaf was wounded and 24 hours later juice was expressed from the upper, unwounded leaf and assayed. The amount of Inhibitor I was measured in 28 F1 plants while the level of Inhibitor II was measured in 27 of the 28 F1 plants. A control group of thirty untransformed tomato plants, var. Better Boy, was also wounded and the amounts of Inhibitors I and II were measured. Inhibitor proteins were not detected in juice expressed from the leaves of six

- 31 -

unwounded antisense plants and six unwounded control plants. Three quarters of the antisense plants (*i.e.*, those inheriting the antisense construct), responded weakly to wounding compared to the control population of untransformed plants (FIGURES 16B AND 16C). Plants not inheriting the construct produced levels of proteinase inhibitors
5 equal to those of the untransformed control plants.

In six of the 28 F1 antisense plants Inhibitor I synthesis in the distal leaf was less than 15% of the mean control level of $97.2 \pm 4.7 \mu\text{g/ml}$ while Inhibitor II synthesis was undetectable in the distal leaf (mean control level of $122.3 \pm 7.2 \mu\text{g/ml}$). Southern blot analysis of the six least responsive F1 plants suggests that
10 these plants inherited two copies of the antisense construct, although this conclusion must be confirmed by self-fertilizing the plants and demonstrating that none of the F2 progeny produce proteinase inhibitors in response to wounding at levels equal to those of the control plants.

These experiments show that expression of antisense prosystemin mRNA in
15 tomato inhibits the systemic induction of proteinase inhibitor syntheses in response to wounding. It is inferred that antisense prosystemin mRNA prevents the efficient production of prosystemin and, hence, of the mobile systemic wound-signal systemin.

Materials and Methods:

A 747 bp fragment of the prosystemin cDNA was excised from pSYS 1 as a
20 BamHI-Hind III fragment. The BamHI site is located in the bluescript polylinker close to the 5'-end of the cDNA, while the Hind III site is within the cDNA at nucleotide 859 as numbered in FIGURE 7. The antisense cDNA fragment thus contained all of the prosystemin mRNA sequence except for the first seven bp of the coding region, all of the 5'-untranslated region and the last 92 bp of the 3'-untranslated region. The cDNA
25 fragment was placed under the control of the constitutive CaMV 35S promoter by cloning it (in the antisense 3' to 5' orientation), into the polylinker of the binary vector pGA643 digested with Bgl II and Hind III. The antisense construct was transformed into *Agrobacterium* strain LBA 4404 and the recombinant bacteria were used to transform tomato var. Better Boy.

30 Tomato seeds, var. Better Boy, were sterilized by soaking for 15 min in a 15% (v/v) solution of Chlorox containing two or three drops of Tween 20. The seeds were washed four times with distilled water then germinated on medium containing: MS salts (4.3 g/L), agarose (6g/L) and thiamine (1 mg/L), pH 5.8. The germinating plants were grown at 28°C with 16 hr days. Eighty percent of the seeds germinated. After
35 7-10 days, when the first true leaves appeared, the cotyledons were removed from the

- 32 -

seedlings and cut into cubes of edge length 0.2-0.5 cm. The tissue cubes were preconditioned on tobacco feeder plates for two days at 25°C in the dark. Tobacco feeder plates were prepared by subculturing tobacco (NT-1) suspension cells in medium containing: MS sales (4.3 g/L), sucrose (30 g/L), inositol (0.1 g/L), thiamine (1 mg/L), 2, 4-D (0.2 mg/L) and KH_2PO_4 (0.18 g/L) at pH 5.8. The cells were incubated for four days at 25°C in the dark. The cells were plated over the same medium including 0.7% agarose, then incubated under the same conditions as before for a further two days. Pieces of tomato cotyledon were placed on Whatman No. 4 filter paper soaked in tobacco feeder plate medium and overlaid onto the tobacco feeder plates. The pieces of preconditioned tissue were punctured with a 20-gauge needle and infected with *Agrobacterium* by soaking them for thirty min in 15 mL of germination medium containing 10^8 cells/mL. The tissue was blotted dry with sterile filter paper and incubated on tobacco feeder plates for a further two days at 25°C in the dark. The tissue pieces were then washed three times in germination medium, the third wash containing 0.5 g/L of Cefotaxime. The tissue pieces were blotted dry with sterile filter paper and placed on shooting medium containing: MS sales (4.3 g/L), thiamine (10 mg/L), nicotinic acid (1 mg/L), pyridoxine (1 mg/L), inositol (100 mg/L), sucrose (30 g/L), BAP (2.5 mg/L), IAA (1 mg/L), cefotaxime (250 mg/L), carbenicillin (500 mg/L), kanamycin (100 mg/L) and 0.7% (w/v) agarose. The explants were transferred after the first three days of culture and weekly thereafter. Once callus growth was observed (after the third subculture) the explants were transferred to shooting medium from which the IAA and BAP had been removed and zeatin (2 mg/L) added. Once the shoots were 2-3 inches tall they were transferred to rooting medium which differed from shooting medium in the BAP, cefotaxime and carbenicillin were absent, vancomycin (0.5 g/L) was added and the concentrations of sucrose (20 g/L), kanamycin (20 mg/L) and IAA (0.05 mg/L) were reduced.

EXAMPLE 11

Effect of growth on Manduca sexta larvae feeding on transgenic plants.

Northern blot analysis of total RNA extracts, collected at different times from undamaged leaves of control (nontransformed) and transgenic tomato plants was performed during feeding experiments with *Manduca sexta* larvae. Plants were approximately 18 inches in height, having 2-3 main stems. Total RNA from each sample was separated by electrophoresis, blotted onto nitrocellulose paper and probed with a nick-translated ^{32}P -prosystemin cDNA. The results are shown in Figure 17.

- 33 -

The accumulation of Inhibitor I and II proteins in undamaged leaves of control and transgenic antisense tomato plants, induced by feeding on *Manduca sexta* larvae was measured. Leaf juice was expressed with a mortar and pestle and assayed by radial immunodiffusion assay (Ryan, 1967). As shown in Figure 18, transgenic antisense tomato plants showed decreased accumulation of both Inhibitor I and II proteins. In Figure 18, A = transgenic antisense plants, C = wild type plants, PI = Inhibitor I and PII = Inhibitor II.

The growth of *Manduca sexta* larvae, while feeding on leaves of control and transgenic antisense tomato plants was measured. Ten first instar larvae were placed randomly on each of six control (wild type) and transgenic plants and removed and weighed. Those larvae feeding on control plants were weighed at 14 days and those feeding on transgenic antisense plants were weighed at 10 and 14 days. As shown in Figure 19, those larvae feeding on transgenic antisense plants showed increased growth weight over those feeding on wild type plants. Values in Figure 19 are the average from three different plant experiments, statistical significance is shown.

Transgenic plants containing a prosystemin gene driven by a CaMV promoter (Example 10, above) were produced. The prosystemin gene was essentially the same as the antisense gene (McGurl *et al.*, *Science* 255, 1570-1573), except the prosystemin cDNA was in the sense orientation. *Manduca sexta* larvae were allowed to feed on leaves of control (wild type) and transgenic sense tomato plants. The control leaves contained little or no Inhibitor I or II. The leaves of the transgenic plant expressing prosystemin contained about 300 μ /g tissue Inhibitor I and about 200 μ /g tissue Inhibitor II. As shown in Figure 20, the weight of larvae feeding on the control leaves increased at a faster rate than the larvae feeding on the transgenic sense leaves.

Those skilled in the art can now appreciate from the foregoing description that the broad teachings of the present invention can be implemented in a variety of forms. Therefore, while this invention has been described in connection with particular examples thereof, the true scope of the invention should not be so limited since other modifications will become apparent to the skilled practitioner upon a study of the specification and following claims.

All publications and applications cited herein are incorporated by reference.

CITATIONS

- 5 1. C.A. Ryan, *Ann. Rev. Phytopathol.* 28, 425 (1990).
2. D.J. Bowles, *Ann. Rev. Biochem.* 59, 873 (1990).
3. M. Chessin and A.E. Zipf, *The Botanical Review* 56, 193 (1990).
- 10 4. D.L. Dreyer and B.C. Campbell, *Plant, Cell and Environ.* 10, 353 (1987).
5. T.R. Green and C.A. Ryan, *Science* 175, 776 (1972).
- 15 6. C.A. Ryan, *TIBS* 3, No. 7, 148 (1978).
7. V.A. Hilder, A.M.R. Gatehouse, S.E. Sheerman, R.F. Barker, D. Boulter, *Nature* 330, 160 (1987).
- 20 8. R. Johnson, J. Narvaez, G. An, C.A. Ryan, *Proc. Natl. Acad. Sci. U.S.A.* 86, 9871 (1989).
9. J.S. Graham, G. Hall, G. Pearce, C.A. Ryan, *Planta* 169, 399 (1986).
- 25 10. J.S. Graham, G. Pearce, J. Merryweather, K. Titani, L. Ericsson, C.A. Ryan, *J. Biol. Chem.* 260, No. 11, 6555 (1985).
11. J.S. Graham, G. Pearce, J. Merryweather, K. Titani, L.H. Ericsson, C.A. Ryan, *J. Biol. Chem.* 260, No. 11, 6561 (1985).
- 30 12. C.A. Ryan, *Plant Physiol.* 43, 1880 (1968).
13. W.E. Brown and C.A. Ryan, *Biochemistry* 23, 3418 (1984).
- 35 14. W.E. Brown, K. Takio, K. Titani, C.A. Ryan, *Biochemistry* 24, 2105 (1985).
15. D. Roby, A. Toppan, M.T. Esquerre-Tugaye, *Physiol. Mol. Pl. Pathol.* 30, 6453 (1987).
- 40 16. H.D. Bradshaw, J.B. Hoflick, T.J. Parsons, H.R.G. Clarke, *Plant Mol. Biol.* 14, 51 (1989).
17. C.A. Ryan and E.E. Fanner, *Annu. Rev. Plant. Physiol. Mol. Bio.* 42, 651 (1991).
- 45 18. E.E. Farmer and C.A. Ryan, *Proc. Natl. Acad. Sci. U.S.A.* 87, 7713 (1990).
- 50 19. H. Pena-Cortes, J.J. Sanchez-Serrano, R. Mertens, L. Willmitzer, S. Prat, *Proc. Natl. Acad. Sci. U.S.A.* 86, 9851 (1989).
20. E. Davies, *Plant, Cell and Environ.* 10, 623 (1987).

- 35 -

21. J.F. Thain, H.M. Doherty, D.J. Bowles, D.C. Wildon, *Plant, Cell and Environ.* 13, 569 (1990).
- 5 22. G. Pearce, D. Strydom, S. Johnson, C.A. Ryan, *Science* 253, 895 (1991).
23. B. McGurl, G. Pearce and C.A. Ryan, *Plant Molecular Biology*, submitted.
- 10 24. H.A. Lutcke *et al.*, *EMBO Journal* 6, 43 (1987).
25. R.B. Harris, *Arch. Biochem. Biophys.* 275, No. 2, 315 (1989).
- 15 26. J. Douglass, O. Civelli and E. Herbert, *Ann. Rev. Biochem.* 53, 665 (1984).
27. L.J. Jung and R.H. Scheffler, *Science* 251, 1330 (1991).
- 20 28. C.A. Ryan, *Anal. Biochem.* 19, 434 (1967).
29. R. Trautman, K.M. Cowan, G.G. Wagner, *Immunochemistry* 8, 901 (1971).
- 25 30. T.P. Hopp and K.R. Woods, *Proc. Nat. Acad. Sci.* 78, 3824 (1981).
31. I. Schechter and A. Berger, *Biochem. Biophys. Res. Commun.* 27, 157 (1967).
- 30 32. S.O. Rogers and A.J. Bendich, *Plant Mol. Biol.* 5, 69 (1985).

- 36 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT: Ryan, C.A.; McGurl, B.F.; Pearce, G.L.
 (ii) TITLE OF INVENTION: "SYSTEMIN"
 (iii) NUMBER OF SEQUENCES: 4
 (iv) CORRESPONDENCE ADDRESS:
 10 (A) ADDRESSEE: Christensen, O'Connor, Johnson and
 Kindness
 (B) STREET: 2800 Pacific First Center, 1420 Fifth Avenue
 (C) CITY: Seattle
 (D) STATE: Washington
 (E) COUNTRY: USA
 15 (F) ZIP: 98101-2347
 (v) COMPUTER READABLE FORM:
 (A) MEDIUM TYPE: Diskette-5.25 inch, 1.2Mb storage
 (B) COMPUTER: IBM PC/386 Compatible
 (C) OPERATING SYSTEM: MS-DOS 4.01
 20 (D) SOFTWARE: Word for Windows-t
 (vi) CURRENT APPLICATION DATA:
 (A) APPLICATION NUMBER: none
 (B) FILING DATE: 03/19/92
 (C) CLASSIFICATION:
 25 (vii) PRIOR APPLICATION DATA:
 (A) APPLICATION NUMBER: none
 (B) FILING DATE: none
 (viii) ATTORNEY/AGENT INFORMATION:
 (A) NAME: Sundsmo, John, S.
 30 (B) REGISTRATION NUMBER: 34,446
 (C) REFERENCE/DOCKET NUMBER: WSUR-1-6171
 (ix) TELECOMMUNICATION INFORMATION
 (A) TELEPHONE: 1-206-682-8100; 1-206-224-0727 (direct)
 (B) TELEFAX: 1-206-224-0779
 35 (C) TELEX: 4938023

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 40 (A) LENGTH: 200 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: polypeptide
 (A) DESCRIPTION: prosystemin #1-#200; systemin #179-196;

Figure 6-7.

- 45 (ix) SEQUENCE DESCRIPTION: SEQ ID NO:1:
 Met Gly Thr Pro Ser Tyr Asp Ile Lys Asn Lys Gly Asp Asp Met Gln
 1 5 10 15
 Glu Glu Pro Lys Val Lys Leu His His Glu Lys Gly Gly Asp Glu Lys
 50 20 25 30
 Glu Lys Ile Ile Glu Lys Glu Thr Pro Ser Gln Asp Ile Asn Asn Lys
 35 40 45

- 37 -

Asp Thr Ile Ser Ser Tyr Val Leu Arg Asp Asp Thr Gln Glu Ile Pro
 50 55 60

5 Lys Met Glu His Glu Glu Gly Gly Tyr Val Lys Glu Lys Ile Val Glu
 65 70 75 80

Lys Glu Thr Ile Ser Gln Tyr Ile Ile Lys Ile Glu Gly Asp Asp Asp
 85 90 95

10 Ala Gln Glu Lys Leu Lys Val Glu Tyr Glu Glu Glu Glu Tyr Glu Lys
 100 105 110

Glu Lys Ile Val Glu Lys Glu Thr Pro Ser Gln Asp Ile Asn Asn Lys
 115 120 125

15 Gly Asp Asp Ala Gln Glu Lys Pro Lys Val Glu His Glu Glu Gly Asp
 130 135 140

20 Asp Lys Glu Thr Pro Ser Gln Asp Ile Ile Lys Met Glu Gly Glu Gly
 145 150 155 160

Ala Leu Glu Ile Thr Lys Val Val Cys Glu Lys Ile Ile Val Arg Glu
 165 170 175

25 Asp Leu Ala Val Gln Ser Lys Pro Pro Ser Lys Arg Asp Pro Pro Lys
 180 185 190

Met Gln Thr Asp Asn Asn Lys Leu
 195 200

30 (2) INFORMATION FOR SEQ ID NO:2:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 951 bases
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: cDNA
 (A) DESCRIPTION: prosystemin cDNA; start ATG at #105
 (ix) SEQUENCE DESCRIPTION: SEQ ID NO:2:

35 AAAATTAAAT TTGATATTG GTTAACTCG ATTTTCCATG AACACCCTTA GTGATGAGTA 60
 TATAAAGCTC AGCTCATGAA GAGTTGAAAT AAAC TAAGAA AACCATGGGA ACTCCTTCAT 120
 ATGATATCAA AAACAAAGGA GATGACATGC AAGAAGAACC AAAGGTGAAA CTTCAACCATG 180
 45 AGAAGGGAGG AGATGAAAAG GAAAAAATAA TTGAAAAGA GACTCCATCC CAAGATATCA 240
 ACAACAAAGA TACCATCTCT TCATATGTTT TAAGAGATGA TACACAAGAA ATACCAAAGA 300
 TGGAACATGA GGAGGGAGGA TATGTAAAGG AGAAAATTGT TGAAAAGGAG ACTATATCCC 360
 50 AATATATCAT CAAGATTGAA GGAGATGATG ATGCACAAGA AAACTAAAG GTTGAGTATG 420
 AGGAGGAAGA ATATGAAAAA GAGAAAATAG TTGAAAAGA GACTCCATCC CAAGATATCA 480

- 38 -

ACAACAAAGG AGATGATGCA CAAGAAAAAC CAAAGGTGGA ACATGAGGAA GGAGATGACA 540
 AAGAGACTCC ATCACAAGAT ATCATCAAGA TGGGAAGGGGA GGGTGCACTA GAAATAACAA 600
 5 AGGTGCTATC TGAGAAAATT ATAGTACGAG AAGATCTTGC TGTTCATCA AAACCTCCAT 660
 CAAAGCGTGA TCCTCCCAA ATGCAAACAG ACAATAATAA ACTCTAGAAA CATCCAAAAA 720
 10 AAATTAATAA ATAAAAAATT ATAATTCAGA ACGATAAAGT AAAAATTCTG AATTGTCTC 780
 CCGTTAGAAA AGTAACTTCA AATAAATATT TGTCTTTCTT TGTATTTTCA AAGTGTAAAT 840
 TGCTTATTGT ACTTTGAGAA GCTTTCTTTA GATTGTTATG TACTTGTATT GCTTCCTTTC 900
 15 TTTTGGCTTA TTTATATAAT ATAAATAAAA AATAAATAAA TATCTAAAGA T 951

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(A) DESCRIPTION: systemin; Figure 3.

(ix) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Ala	Val	Gln	Ser	Lys	Pro	Pro	Ser	Lys	Arg	Asp	Pro	Pro	Lys	Met	Gln	Thr	Asp
1				5				10					15				

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 4526 bases

(B) TYPE: nucleic acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(A) DESCRIPTION: systemin genomic DNA; Figures 8A-8C.

(ix) SEQUENCE DESCRIPTION: SEQ ID NO:4:

AAAATTAAAT TTGATATTTG GTTAACTCG ATTTTCCATG AACACCCTTA GTGATGAGTA 60
 TATAAAGCTC AGCTCATGAA GAGTTGAAAT AAACAAAGAA AACCATGGGA ACTCCTTCAT 120
 40 ATGATATCAA AAACAAAGGT ATCATTTCTT TATATGCCTA AGTATATATT TATTTATATA 180
 TTTTGTAAGT AAAATTTTAT ATTAAATCA ACAAGTGAGA GTTTAACAAA AATCATATTA 240
 45 AAGAAAAAAA ATATATTAAT TTTTAATCAT GGTATTATCC TCCAGGAGAT GACATGCAAG 300
 AAGAACCAA GGTGAACTT CACCATGAGA AGGTAACCTT AGTTTCTCCT TTTTCTTTTT 360
 TCAACTTCTT TATATATTAT TTTTGTAAT TTTTTATAT TATAATGTTT AAATGGTCTC 420
 50 ATTTTCTAAT TAATAATGTG TCTGAATCGC CATGTTATTT ATGTTAGATT TAATACATTA 480
 ATAACATTGT TAGTAAATGT TAGAATACTG ACTCCCAAAT TCGCTTAAGG AACAAGTATA 540

- 39 -

5 TTTTCATGTGT TTCTTTGCAG ATAACAATAA TTATGTTTTG TAAAGCAAAT AAAATAATAA 600
 CATAATATTT TATCGTAGAA AACTCCAAC TATTATTTAG ATATTTAGAT TATGATTTGC 660
 10 TTTAATTATA CTTTTTTAAA CATGATAAAT TATTTCTGTT AGACATTTTC GATTCATTTT 720
 TTTTTTTACA AAAATTGTAT TTGCTCTCAA ACGTTTACTA GTTAGTTAAG TTAACATATAC 780
 AAAATATGTC ATCTCATTTG ATTATATACA TCAGGCTCAA TTAAACATA TTGGAGATAT 840
 15 GGAGATTTTA CGATTCATTA AACTAATGT GTATAGTTAG AAAATGTGAA ATATTTCAA 900
 TGGTTAACTT TTCTGTATAA TTGACATTTG AACTATATG TTTAATTATA ACAACCGTA 960
 ATCAAATGTT CAAATAAAAT TGAATGACAA TAGGTATAAG GAGCTATCAA TATATTAGCT 1020
 CTTCTTGATT CAACTTATTT ACCGTTATAA TTAAATAATG ACTCGTTAAT TGATTTAATT 1080
 20 TTTTTACTCA CGTGAAATGA TTTAATCAAC TCATTTATCA CCCTTATTTA CGACTCATGT 1140
 AGAATAATGT TCTTTATACT TGTATACAAT TTACTCGGAT ATTTTTTTTA AATTTTTTTT 1200
 TATGTTTAAT TAAATACTAT TAAATGAAG AAATATTATT TATAATTGAA GAATATTGAA 1260
 25 TTTTTTTTCC ATCAAATTT ACAGGGAGGA GATGAAAAGG AAAAAATAAT TGAAAAAGAG 1320
 ACTCCATCCC AAGATATCAA CAACAAAGAT ACCATCTCTT CATATGTTTT AAGTATTTAA 1380
 TTTTTTTCAA TCTTTTTTTT TTCTCATCTT CTTATTTTAA TCATCTAAAA GAAATTATTA 1440
 30 TTATGTTTTT TTTTAACTTT AATTATAATA TTATCCAGCA GGAGATGATA CACAAGAAAT 1500
 ACCAAAGATG GAACATGAGG AGGTAACAT ATATTTCAAT TTATTTACTA ATTTATAAAT 1560
 35 AATGACTTAT TCATTGATTC AATTTATTTT AATTCGTTTG AATCAAAC AAGGTTACCA 1620
 TATTATCACC CCACTCCCTC CACTACTCAT TTAAATGAT GGTTCGATAC TTTGCATGCA 1680
 ATTTTGTTTA TTCATAAGTC ATTTATTTTT CAAAAATTT ATGTTTCAGT AAACGTTTGC 1740
 40 ATACATTTTG TTTATACATA ATTCATCTAT TTCTTTTAAA ATTTTATGTT CAGTTAAACG 1800
 ATTGCATACA TTTTGTTTAT ACATAAGTCA TCTATTTTTT TAAAAAAA TTATGTTTACG 1860
 45 TTAACGTTTG CATACAATTT TGTTTCATACA TAATTCATCT ATTTTTTTTAA AATTTTATGA 1920
 TCAGTTAAAC TTTTATATAC AATTTTGCTC GTACATAAGT CATCTATTTT TTTAAAATTT 1980
 50 TATGTTTCACT TAAACGTAAT AAATAAAATT AGACTGTGGA AATATTATTT ATTATTAAAG 2040
 GATATTACAG GGAGGATATG TAAAGGAGAA AATTGTTGAA AAGGAGACTA TATCCCAATA 2100
 TATCATCAAG ATTGAAGGTA TAATCTATTT ATATGTGTCT AAATATTTAA TTTTATTTTT 2160

- 40 -

ATTTTTCAGA TTTTITAGTA AGGGATTTTT TTATTTTTTT TCAAAAAATG TGAATCATT 2220
TCAAGAAGTT AATATTATTT TTGGTAACTT TAATCTTGAT ATATTATTCT CCAGGAGATG 2280
5 ATGATGCACA AGAAAACTA AAGGTTGAGT ATGAGGAGGT AACTTTAATT TCTTCTTTGA 2340
CTTTTATTT ATTATTTTGT TATATTTTAC TGTCTATTTA TTTTCATATTC ACAAATTATA 2400
TTTATCACAT ATATATGCTT TTATTTTCTT CAAAATTACA GGAAGAATAT GAAAAAGAGA 2460
10 AAATAGTTGA AAAAGAGACT CCATCCCAAG ATATCAACAA CAAAGGTATA TATCATATCT 2520
TCATATGCCT AAGATTTTAT TTTCTTCTTA TTTTTCATAT TATTTTTATT TAACTAAATT 2580
TAGTATGAAA CCTTTTTTTTT TTTAAAAAAA TCATCTTAAA TAAAATATTA TTTTGGGTG 2640
15 ACTCAAATCA TTGACCTTAT ATATTCTCCA GGAGATGATG CACAAGAAAA ACCAAAGGTG 2700
GAACATGAGG TAACTACTTA TATTTTTCTC TCTCTTTATT ACATAAAATC ACATTAGTTA 2760
20 TATGATAATT GGCTATGCTA ATAATAAAAA AACAAATTAAT ATATTTATAG GAATTTAAAC 2820
AGGGTGGAGT GTCCATGATC TTTATTTTTTA TCTTGTAAG TTAATAAGAC TATTTCCAAA 2880
TAGACCTTTA GTTTGAGCAA AATCTATCAG AAAATACGAT AATAAAGAAG TCACGCTGAA 2940
25 AATAAAATAT TAATTTTGTG ACGTGAAAGC AATATCAAGA GCCCCGTCAA TTTGTTGTAT 3000
TATGTCAGAT GCAACATCCT TCTTTCTTCT CGTGAAGTAT AGGAGCGCTT AGCACACATC 3060
30 TCAACATAAT GCGCGATAAT AACGTTTTAA TGGTGAATCT ATCGGTATCA TAACAATAGT 3120
ATACAACCTT AAACCTAATG ATCGTCTAGC TAGTAATCTT TCAAATGAG GGACCCTAAT 3180
TACTGACAAA ATTTGTGTCT AACATAACTT ATGTACCATA ACAATAATAT ATCTTGTGTA 3240
35 ATTTATGAGT GAAGGTAGGG TTTGAAATTA AACATAATCA ATAAAATTGG ACAAAGAAGA 3300
TATTATTTAT TAATTGAAAG ATATTAATAG TTTTCTTCTT CAAAATTACA GGAAGGAGAT 3360
40 GACAAAGAGA CTCCATCACA AGATATCATC AAGATGGAAG GTATCAATCT ATTTATATTT 3420
TTTATAAGTA TTTTCTTCTT ACAATTTTTT TATTTCTTTT GGTATATATG AAACATTTTT 3480
TTTAAACCAT CTTTAAAAAA AAATAATACT TATGTATAAC TATAATCATG ATATTATCAT 3540
CCAGGGGAGG GTGCACTAGA AATAACAAAG GTGGTATGTG AGGTAACTAA ATTTCTTCTT 3600
CCAATTTTTC TATACATTAT GTTTGTATTT TTTTCTTGGG ATTCATTCTG ACTTTCTTCT 3660
50 ATAGAAAGTC TTGCTATCTA TATACGATTA AAATTATATT GAGTTTACGA TAAAAATATA 3720
TTTAAACAAT TCTTTTTTTT ATTTTCATATC TAACTATTG AAAATGTGTC TGCCCTCGTA 3780

- 41 -

ACCTCGGTAC AAAGCCAACT AGAACCACAT TTAAATGAT TAAAAAATC TTTTGAAAGT 3840
GTGAGAAATA CGCTGAACT ATCGCTTATT ATTTTATTTT TACGTATATG CAATAGACAA 3900
5 TATTGAATCC TCTTCTATTT ATTCGTATGT TTACTTCCTC ACATATCAAA TCTCTTAGTA 3960
AAAATTCTGA CTCACCACT GTATATATCT TTTTATTTTG ATTTTGTATT GCATTCATT 4020
10 TGTTTAGTTA TAATAACTAA TAAGCGTCTT TTATTTTATT TATAGCATGA TGCTACTATT 4080
TTTTGGACAC TACAAGGAGC ATACAATTCA AATCTCAAAC TTTTTATAT TTTTTTCTA 4140
TATTTTTTAT TATAAAGGA TATTAATTC TTTTTCTTT CAAATACAGA AAATTATAGT 4200
15 ACGAGAAGAT CTGCTGTTT AATCAAAACC TCCATCAAAG CGTGATCCTC CCAAAATGCA 4260
AACAGACAAT AATAAATCT AGAAACATCC AAAAAAATT AATAAATAAA AAATTATAAT 4320
20 TCAGAACGAT AAAGTAAAA TTCTGAATTT GTCTCCCGTT AGAAAAGTAA CTTCAAATAA 4380
ATATTTGTCT TTCTTTGTAT TTTCAAAGTG TAATTGGTT ATTGTACTTT GAGAAGCTTT 4940
CTTTAGATTG TTATGTACTT GTATTGCTTC CTTTCTTTTG GCTTATTTAT ATAATATAAA 4500
25 TAAAAAATAA ATAAATATCT AAAGAT 4526

- 42 -

WHAT IS CLAIMED IS:

1. A nucleic acid comprising a nucleotide sequence substantially as shown in SEQ. ID. NO. 2.
5
2. A nucleic acid comprising a nucleotide sequence substantially complementary to the sequence of Claim 1.
3. The nucleic acid of Claim 1, further comprising the nucleotide sequence
10 of residues number 639 through 692 of SEQ. ID. NO. 2.
4. The nucleic acid of Claim 1, further comprising a synthetic oligonucleotide.
- 15 5. A recombinant nucleic acid vector comprising the nucleic acid sequence of Claim 1 operably linked to a promoter nucleotide sequence.
6. A polypeptide encoded by the nucleic acid of Claim 1.
- 20 7. The nucleic acid of Claim 1 capable of encoding an antisense RNA.
8. A recombinant nucleic acid vector comprising the nucleic acid sequence of Claim 2 operably linked to a promoter nucleotide sequence.
- 25 9. A nucleic acid comprising a nucleotide sequence substantially as shown in SEQ. ID. NO. 4.
10. A nucleic acid comprising a nucleotide sequence substantially complementary to the sequence of Claim 9.
30
11. The nucleic acid of Claim 9, further comprising a synthetic oligonucleotide.
12. A polypeptide having an amino acid sequence substantially as shown
35 in SEQ. ID. NO. 1.

- 43 -

13. A polypeptide having an amino acid sequence substantially as shown in SEQ. ID. NO. 3.

14. A polypeptide comprising the amino acid sequence
5 R₁R₁QR₁R₂PPR₁R₂R₂R₁PPR₂R₁QR₁R₁, wherein R₁ is any amino acid, R₂ is lysine or arginine, Q is glutamine, and P is proline.

15. The polypeptide of Claim 14, wherein the polypeptide comprises the amino acid sequence NH₃-AVQSKPPSKRDPPKMQTD-COO⁻.
10

16. A process for inhibiting synthesis of a defense protein in a plant comprising introducing the nucleic acid sequence of Claim 2 into a plant cell and growing the cell into a plant.

17. A transgenic plant produced by the process of Claim 16, or by breeding the plant of Claim 16 with a second plant.
15

18. A transgenic plant comprising a recombinant nucleic acid that encodes systemin or prosystemin.
20

19. The transgenic plant of Claim 18, further comprising a plant selected from the group consisting essentially of tomato, potato, tobacco, alfalfa, rape, soybean, corn, rice, wheat, barley, cotton, poplar tree, pine and fir tree.

20. A method of enhancing the defense mechanism of a plant comprising the step of introducing to the plant a nucleic acid coding for a polypeptide hormone which stimulates the production of a defense protein in the plant.
25

21. The method of Claim 20, wherein the polypeptide hormone is systemin.
30

22. Substantially pure systemin transcript.

23. The DNA template sequence for the transcript of Claim 22.

- 44 -

24. The DNA sequence substantially complementary to the template sequence of Claim 22.

5

25. The amino acid sequence translated from the transcript of Claim 22.

26. The transcript of Claim 22, wherein the transcript is not of *in vivo* origin.


10 27. A method of increasing the level of polypeptide hormone in a plant comprising the step of introducing a recombinant nucleic acid coding for the polypeptide hormone.

28. The method of Claim 27, wherein the polypeptide hormone is systemin.

1/28

Inhibitor I ($\mu\text{g/g}$ tissue) (---x--)

FIGURE 1



SUBSTITUTE SHEET

2/28




FIGURE 2

3/28



FIGURE 3

4/28




FIGURE 4

SUBSTITUTE SHEET

5/28

Fig. 5A.




Fig. 5B.

SUBSTITUTE SHEET

6/28

Met	Gly	Thr	Pro	Ser	Tyr	Asp	Ile	Lys	Asn	Lys	Gly	Asp	Asp	Met	Gln	1	5	10	15
Glu	Glu	Pro	Lys	Val	Lys	Leu	His	His	Glu	Lys	Gly	Gly	Asp	Glu	Lys	20	25	30	
Glu	Lys	Ile	Ile	Glu	Lys	Glu	Thr	Pro	Ser	Gln	Asp	Ile	Asn	Asn	Lys	35	40	45	
Asp	Thr	Ile	Ser	Ser	Tyr	Val	Leu	Arg	Asp	Asp	Thr	Gln	Glu	Ile	Pro	50	55	60	
Lys	Met	Glu	His	Glu	Glu	Gly	Gly	Tyr	Val	Lys	Glu	Lys	Ile	Val	Glu	65	70	75	80
Lys	Glu	Thr	Ile	Ser	Gln	Tyr	Ile	Ile	Lys	Ile	Glu	Gly	Asp	Asp	Asp	85	90	95	
Ala	Gln	Glu	Lys	Leu	Lys	Val	Glu	Tyr	Glu	Glu	Glu	Glu	Tyr	Glu	Lys	100	105	110	
Glu	Lys	Ile	Val	Glu	Lys	Glu	Thr	Pro	Ser	Gln	Asp	Ile	Asn	Asn	Lys	115	120	125	
Gly	Asp	Asp	Ala	Gln	Glu	Lys	Pro	Lys	Val	Glu	His	Glu	Glu	Gly	Asp	130	135	140	
Asp	Lys	Glu	Thr	Pro	Ser	Gln	Asp	Ile	Ile	Lys	Met	Glu	Gly	Glu	Gly	145	150	155	160
Ala	Leu	Glu	Ile	Thr	Lys	Val	Val	Cys	Glu	Lys	Ile	Ile	Val	Arg	Glu	165	170	175	
Asp	Leu	Ala	Val	Gln	Ser	Lys	Pro	Pro	Ser	Lys	Arg	Asp	Pro	Pro	Lys	180	185	190	
Met	Gln	Thr	Asp	Asn	Asn	Lys	Leu	195	200										

FIGURE 6

FIGURE 7A

8/28

420 GAGGAGGAAGAAATATGAAAAAGAGAAAAATAGTTGAAAAAGAGACTCCATCCCAAGATATC 470
 106 GluGluGluGluTyrGluLysGluLysIleValGluLysGluThrProSerGlnAspIle 125
 490 510 530
 480 AACAAACAAGGAGATGATGCACAAAGAAAAACCAAGGTGGACATGAGGAAGGAGATGAC 539
 126 AsnAsnLysGlyAspAlaGlnGluLysProLysValGluHisGluGluGlyAspAsp 145
 550 570 590
 540 AAAGAGACTCCATCACAAAGATATCATCAAGATGGAAGGGAGGTGCACTAGAAATAACA 599
 146 LysGluThrProSerGlnAspIleIleLysMetGluGlyAlaLeuGluIleThr 165
 610 630 650
 600 AAGGTGGTATGTGAGAAAAATTATAGTACGAGAAGATCTTGCTGTTCAATCAAAACCTCCA 659
 166 LysValValCysGluLysIleIleValArgGluAspLeuAlaValGlnSerLysProPro 185
 670 690 710
 660 TCAAAAGCGTGATCCTCCCAAAATGCACAAACAGACAAATAATAACTCTAGAAACATCCAAA 719
 186 SerLysArgAspProLysMetGlnThrAspAsnAsnLysLeu
 730 750 770
 720 AAAATTAAATAAATAAATAATTAATAATTCAGAACGATAAAAGTAAAAAATTCGAATTGTCT 779
 790 810 830
 780 CCCGTTAGAAAAGTAACCTTCAAAATAAATAATTTGCTCTTTCTTTGTATTTTCAAAAGTGAAT 839
 850 870 890
 840 TTGGTTATTGTACTTTTGAGAAAGCTTTCTTTAGATTGTTATGTACTTGTATTGCTTCCTTT 899
 910 930 950
 900 CTTTGGCTTATTATATAATAATAATAAATAAATAAATAAATAAATAAATAAATAAAGAT 951

FIGURE 7B

SUBSTITUTE SHEET

FIGURE 8A1

1100 GAGCTATCAAATATATTAGCTCTTCTTGATTCAACTTATTACCGTTATAATTAAATGAATGACTCGTTAAATGATTTAAATTTTTTACTCAGTGAAATGA
1200 TTTAATCAACTCAATTTATCACCCTTATTACCGACTCATGTAGAAATAATGTTCTTTTATACITGTATACAAATTTACTCGGAATTTTTTTTTTAAATTTTTTTT
1300 TATGTTTTAAATTAAATAGAGAAAATATTATTATAATTGAAGAATATTGAATTTTTTTTCCATCAAAAATTTACAGGGAGGAGATGAAAAGG
1400 AAAAATAATTGAAAAAGAGACTCCATCCCAAGATATCAACAACAAGATACCATCTCTTCATATGTTTTAAAGTATTTAAATTTTTTTTCAATCTTTTTTTT
1500 TTCTCATCTTCTTATTTTAATCATCTAAAGAAATTTATTATTATGTTTTTTTAACTTTTAAATTAATAATATATCCAGCAGGAGATGATACACAAGAAAT
1600 ACCAAAGATCGAACAATGAGGAGGTAACATAATATTTTCAATTTATTACTAATTTTATAAAATGAATGACTTAATTCATTCGATTCGATTTTAAATTCGTTTG
1700 AAATCAAACTAAGGTTACCATAATTATCACCCTCAGTCCCTCCACTACTCAITTTAAATGAATGTTGATACITTTGCATGCAATTTTGTTTATTCATAAGTC
1800 ATTTATTTTCAAAAATTTTATGTTTCAGTTAAACGTTTGCATACATTTTGTGTTTATACATAATTCATCTAATTTCTTTTTTAAATTTTATGTTCAGTTAAACG
1900 ATTGCATACATTTTGTTCATACATAAGTCATCTAATTTTTTAAAAAAAATTTATGTTTCAGTTAACGTTTGCATACAAATTTTGTTCATACATAATTCATCT

FIGURE 8A2




FIGURE 8B1

2901 AATCTATCAGAAAAACGAATAAAGAACTCAGCGCTGAAATAAAAATATTAATTTTGTGACGTGAAAGCAATATCAAGAGCCCGTCAATTTGTTGTAT
 3001 TATGTCAGATGCAACATCCCTCTTTCTTCTGGAAGTATAGCAGGGCTTAGCACACATCTCAACATAAAGCGGATAAIAACGTTTAAATGGTGAATCT
 3101 ATCGGTATCATAACAATAGIATACAACTTTAAACCTAATGATCGCTAGCTAGTAATCTTTCAAAATGAGGGACCCTAAATTACTGACAAAATTTGTGTCCT
 3201 AACATAACTIATGTACCATAACAATAATATCTTGTGTAAATTATGAGTGAAGGTAGGGTTTGAAATTTAAACATAATCAATAAAATTTGACAAAGAAGA
 3301 TATTATTTAATTAAGGATATTAATAGTTTTTTTCTTCAAAATTACAGGAAGGAGATGACAAAGAGACTCCATCACAAAGATATCATCAAGATGGAAG
 3401 GTATCAATCTATTATATTTTATAAGIATTTTTTCTTACAATTTTTTATTTCTTTGGTATATATGAAACTATTTTTTTTAAACCACTCTTTAAAAAA
 3501 AAATAATACIATAGIATAACIATAATCATGATATATATCAATCCAGGGGAGGTGCACACTAGAAATAACAAAGGTGGTATGTGAGTAACTAAATTTCTTCTT
 3601 CCAATTTTCTATACATTAIGTTTGATTTTTTTTTTGGATTCAATTCGAACTTTCTTCGATAGAAAGCTTGGCTATCTATATACGATTAAATTTATAT
 3701 GAGTTTACGATAAAAAATATATTTAAACAATCTTTTTTAAATTCATATCTAAACTATTGAAATGTGTCTGCCCTCGTAACTCGGTACAAAGCCCAACT
 3801 AGAACCCACATTTTAAATGATTAATAAAAAATCTTTTGAAAGGTGAGAAATACCGCTGAACIATCGCTTATTTATTTTACGTATATGCAATAGACAA
 3901 TATTGAATCCCTCTCTATTATTTCGTAIGTTTACTTCCICACATATCAAAATCTCTTAGTAAATTTCTGACTTCACCACIATATATCTTTTTTATTG
 4000

12/28

Exon 9

Exon 10

FIGURE 8B2



SUBSTITUTE SHEET

SUBSTITUTE SHEET

FIGURE 8C


14/28

Fi . . . A .



Fi . . . B .

Fi . . . C .



1 kb

FIGURE 10

16/28

Ex3 GgAGgAgAIG gAAgGAgAA AATAgTTGAA AAgAGAGAGTc CATCCCAAGA TATCAACAGC AAGATAcCCc tctcttcata tgtttttttag
Ex7 GgAGAtAIG gAAAgAgAA AATAgTTGAA AAgAGAGAGTc CATCCCAAGA TATCAACAGC AAG.....
Ex5 GgAGgAtAIG LAAAgGAgAA AATtG. JAA AAgGAGAGTc TATCCCAATA TATCATCAAG AttGAgg....
Ex99A AggAGgTCAG AAgAGAGAGTc CATCCCAAGA TATCATCAAG AtgGAgg....
Ex1 AtgGgAgAGTc CtTcAtATGA TATCAAAAGC AAG.....
Cun G-AG-A-AIG -AAA-GA-AA AATAgTTGAA AAgAGAGAGTc CATCCCAAGA TATCAACAGC AAGA-----

FIGURE 11A

```
Ex4  ....AGATGA TGCACAAGAA ATACCAAAGG TCGAACATGA GgAg.....
Ex8  ...GACA1CA TGCACAAGAA AAACCAAAGC TCGAACATGA C.....
Ex6  ggggAGATGA TGCACAAGAA AAACCAAAGC TCGAgATGA GgAg.....
Ex10  ...GgGAgGg TGCACtAGAA AtAaCAAAGC TCGtAtgTGA C.....
Ex2  ...CACATCA catgCAAGAA gAACCAAAGC TCaAACtTca ccAtgggggg
Ctrl ---GAGA1CA TGCACAAGAA AAACCAAAGC TCGAACATGA C-A-----
```

FIGURE 11B

SUBSTITUTE SHEET

	31		48
Rep A	Glu lys glu lys ile <u>ile</u> glu lys glu thr pro ser gln asp ile asn asn lys		
	111		128
Rep 2A	Glu lys glu lys ile <u>val</u> glu lys glu thr pro ser gln asp ile asn asn lys		
	58		71
Rep B	Asp asp thr gln glu <u>ile</u> pro lys <u>met</u> glu his glu glu gly		
	130		143
Rep 2B	Asp asp <u>ala</u> gln glu lys pro lys <u>val</u> glu his glu glu gly		
	81		93
Rep C	Lys glu thr <u>ile</u> ser gln tyr ile ile lys <u>ile</u> glu gly		
	146		158
Rep 2C	Lys glu thr <u>pro</u> ser gln <u>asp</u> ile ile lys <u>met</u> glu gly		

FIGURE 12

SUBSTITUTE SHEET




FIGURE 13A




FIGURE 13B

Exon 3 AAGATACCATCTCTTCATATGTTTTAAG

Exon 7 AAGATATCATATCTTCATATGCCTAAG
 $\swarrow \searrow$
 GTAT

FIGURE 14




Fig. 15A.

Rt St Pt Le Se Pe Sm Pi




Fig. 15B.

Kb 1 2

1.7 ▶

1.0 ▶

Fig. 16 A.

24/28




FIGURE 16B




FIGURE 16C

25/28

CONTROL ANTISENSE
0 6 12 0 6 12 days



Fi .17.

26/28




FIGURE 18

SUBSTITUTE SHEET




FIGURE 19

SUBSTITUTE SHEET




FIGURE 20

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/02428

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 68, 70, 69.1, 91, 172.1, 320.1; 530/379, 380, 381, 382; 536/27.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CA, APS, DIALOG AND ONLINE SEQUENCE SEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	SCIENCE, VOLUME 255, ISSUED 20 MARCH 1992, B. MCGURL ET AL., "STRUCTURE, EXPRESSION, AND ANTISENSE INHIBITION OF THE SYSTEMIN PRECURSOR GENE", PAGES 1570-1573, ENTIRE DOCUMENT.	1-28
Y,P	JOURNAL OF PROTEIN CHEMISTRY, VOLUME 11, NO. 3, ISSUED JUNE 1992, D. J. RUSSELL ET AL., "PROTON NMR ASSIGNMENTS OF SYSTEMIN", PAGES 265-274, ENTIRE DOCUMENT.	1-28
Y	SCIENCE, VOLUME 253, ISSUED 23 AUGUST 1991, G. PEARCE ET AL., "A POLYPEPTIDE FROM TOMATO LEAVES INDUCES WOUND-INDUCIBLE PROTEINASE INHIBITOR PROTEINS", PAGES 895-898, ENTIRE DOCUMENT.	1-28

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 30 APRIL 1993	Date of mailing of the international search report 17 MAY 1993
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer GIAN WANG
Facsimile No. NOT APPLICABLE	Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/02428

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	PROC. NATL. ACAD. SCI. USA, VOLUME 86, ISSUED DECEMBER 1989, R. JOHNSON ET AL., "EXPRESSION OF PROTEINASE INHIBITORS I AND II IN TRANSGENIC TOBACCO PLANTS: EFFECTS ON NATURAL DEFENSE AGAINST MANDUCA SEXTA LARVAE", PAGES 9871-9875, ENTIRE DOCUMENT.	1-28
Y	US, A, 4,394,443 (WEISSMAN ET AL.) 19 JULY 1983, ENTIRE DOCUMENT.	1-28

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/02428

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (5):

C07H 15/12; C07K 3/00, 13/00, 15/00, 17/00; A61K 35/14, 35/78, 35/80, 37/04; C12Q 1/68; C12N 15/00; C12P 21/00, 21/02, 21/06, 19/34.

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/6, 68, 70, 69.1, 91, 172.1, 320.1; 530/379, 380, 381, 382; 536/27.

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)